

Les Cahiers de la Chaire / N°58

A new class of problems in the calculus of variations

Ivar Ekeland, Yiming Long & Qinglong Zhou



CHAIRE

Finance & Développement Durable

A new class of problems in the calculus of variations

Ivar Ekeland^{1*}, Yiming Long^{2†}, Qinglong Zhou^{2‡}

¹ CEREMADE and Institut de Finance
Université de Paris-IX, Dauphine, Paris, France

² Chern Institute of Mathematics and LPMC
Nankai University, Tianjin 300071, China

June 19, 2013

Dedicated to Professor Alain Chenciner on his 70th birthday

1 Introduction

In economic theory, and in optimal control, it has been customary to discount future gains at a constant rate $\delta > 0$. If an individual with utility function $u(c)$ has the choice between several streams of consumption $c(t)$, $0 \leq t$, he or she will choose the one which maximises the present value, given by:

$$\int_0^{\infty} u(c(t)) e^{-\delta t} dt \quad (1)$$

That future gains should be discounted is well grounded in fact. On the one hand, humans prefer to enjoy goods sooner than later (and to suffer bads later than sooner), as every child-rearing parent knows. On the other hand, it is also a reflection of our own mortality: 10 years from now, I may simply no longer be around to enjoy whatever I have been promised. These are two good reasons why people are willing to pay a little bit extra to hasten the delivery date, or will require compensation for postponement, which is the essence of discounting.

On the other hand, there is no reason why the discount rate should be constant, i.e. why the discount factor should be an exponential $e^{-\delta t}$. The practice probably arises from the compound interest formula $\lim_{\varepsilon \rightarrow 0} (1 - \varepsilon \delta)^{t/\varepsilon} = e^{-\delta t}$, when a constant interest rate δ is assumed, but even in finance, interest rates vary with the horizon: long-term rates can be widely different from short-term ones. As for economics, there is by now a huge amount of evidence that individuals use higher discount rates for the near future than for the long-term (see [18] for a review up to 2002). There

*E-mail: ekeland@math.ubc.ca

†Partially supported by NNSF, MCME, RFDP, LPMC of MOE of China, and Nankai University. E-mail: longym@nankai.edu.cn

‡Partially supported by the Chern Institute of Mathematics, Nankai University, CEREMADE of Université Paris Dauphine, and the IHES. E-mail: zhou.qinglong.1985@gmail.com

is also an aggregation problem: in a society where individuals use constant (but different) discount rates, the collective discount rate may be non-constant (see [14]). So the present value formula (1) should be replaced by the more general one:

$$\int_0^{\infty} u(c(t)) h(t) dt \quad (2)$$

where h is a decreasing function, with $h(0) = 1$.

But then a new problem arises, which is now well recognized in economic theory, but to our knowledge has not yet received the attention it deserves in control theory. It is the problem of *time-inconsistency*, which runs as follows. Suppose the decision-maker has the choice between two streams of consumption $c_1(t)$ and $c_2(t)$, starting at time $T > 0$. At time $t = 0$, he or she finds $c_1(t)$ yields the highest present value:

$$\int_T^{\infty} u(c_1(t)) h(t) dt > \int_T^{\infty} u(c_2(t)) h(t) dt. \quad (3)$$

He or she then chooses $c_1(t)$. When time T is reached, the present values are now:

$$\int_T^{\infty} u(c_1(t)) h(t-T) dt \quad \text{and} \quad \int_T^{\infty} u(c_2(t)) h(t-T) dt \quad (4)$$

If $h(t) = e^{-\delta t}$, then the ordering found at time $t = 0$ will persist at time $t = T$. Indeed:

$$\int_T^{\infty} u(c(t)) e^{-\delta(t-T)} dt = e^{\delta T} \int_T^{\infty} u(c(t)) e^{-\delta t} dt$$

so that the two terms in (4) are proportional to the two terms in (3). However, this is a peculiarity of the exponential function, and it is not to be expected with more general discount rates. The decision-maker then faces a basic rationality problem: what should he or she do? To be more specific, assume the state $k(t)$ is related to the control $c(t)$ by the dynamics:

$$\frac{dk}{dt} = f(k) - c(t), \quad k(0) = k_0 \quad (5)$$

$$c(t) \geq 0, \quad k(t) \geq 0 \quad (6)$$

and the decision-maker is interested in maximising (2). How should he or she behave?

In the exponential case, when $h(t) = e^{-\delta t}$, the answer is to pick the optimal solution: if it is optimal at time $t = 0$, it will still be optimal at all times $T > 0$ (this, by the way, is the content of the dynamic programming principle). But in the non-exponential case, the notion of optimality changes with time: each observer, from time $t = 0$ on, has his or her own optimal solution. No one agrees on what the optimal solution is, so optimality no longer provides an answer to the decision-making process, and one must look for other concepts to describe rational behaviour.

A clear requirement for rationality is that any strategy put forward be implementable. Suppose a Markov strategy $c = \sigma(k)$ is put forward at time $t = 0$. If it is to be followed at all later times $t > 0$, then it must be the case that the decision-maker at that time finds no incentive to deviate. More precisely, if he/she assumes that at all later times the strategy (closed-loop feedback) $c = \sigma(k)$ will be applied, then he/she should find it in his/her interest to apply σ as well. In other words, σ

should be a subgame-perfect Nash equilibrium of the leader-follower game played by the successive decision-makers. This idea has been introduced by Phelps ([29], [30]) in models with discrete time (see [24] and [27] for further developments), and adapted by Karp ([22], [23]), and by Ekeland and Lazrak ([12], [13], [14], [16]) to the case of continuous time.

In this paper, we will follow the approach by Ekeland and Lazrak. It consists of introducing a value function $V(k)$, which is very similar to the value function in optimal control, and of showing that it satisfies a functional-differential equation which is reminiscent of the Hamilton-Jacobi-Bellman (HJB) of optimal control. Conversely, any solution of that equation with suitable boundary conditions will give us an equilibrium strategy.

In the work by Ekeland and Lazrak, this approach was applied to (2), with $h(t) = \alpha \exp(-r_1 t) + (1 - \alpha) \exp(-r_2 t)$, and it was showed that the corresponding problem had a continuum of equilibrium strategies. In the present paper, in view of applications to economics, and of the mathematical interest, we aim to extend the analysis to the more general case:

$$(1 - \alpha) \int_0^\infty e^{-r_1 t} u(c(t), k(t)) dt + \alpha \int_0^\infty e^{-r_2 t} U(c(t), k(t)) dt \quad (7)$$

As a by-product of our analysis, we will treat the problem:

$$(1 - \alpha) \delta \int_0^\infty e^{-\delta t} u(c(t), k(t)) dt + \alpha \lim_{t \rightarrow \infty} U(k(t), c(t)) \quad (8)$$

which was introduced by Chichilnisky (see [10], [11]) to model sustainable development. Note that, if $u(c, k) \geq 0$ and:

$$\sup \{U(c, k) \mid c > 0, k > 0, c = f(k)\} = \infty$$

then maximising (8) under the dynamics (5), (6) leads to the value $+\infty$, so that optimisation is clearly not an answer to the problem. Instead, we find equilibrium strategies. To our knowledge, this is an entirely new result. We show that there is a continuum of such strategies. More precisely, there is a continuum of points k_∞ which can be realized as the long-term level of capital by an equilibrium strategy. This support, however, is one-sided, that is, k_∞ can be reached only from the initial level of capitals k_0 lying on its left (or from its right). To our knowledge, this is the first time such strategies have been identified.

The structure of the paper is as follows. In the next section, we consider the problem of maximising

$$\int_0^\infty e^{-\delta t} u(c(t), k(t)) dt \quad (9)$$

under the dynamics (5), (6), and we show that it has a solution. On the way, we introduce the corresponding HJB equation, and we show that it has a C^2 solution. Next, we define equilibrium strategies. With each such strategy we associate a value function $V(k)$, and we show that it satisfies an integro-differential equation which generalizes the HJB equation, and we prove a verification theorem: any solution of this equation with suitable boundary conditions gives an equilibrium strategy. We show that the trajectories satisfy an integro-differential equation which generalizes the classical Euler-Lagrange equations, and we connect the Ekeland-Lazrak approach with the Karp approach.

In section 4, we apply the theory to problem (7), and show that it has a continuum of equilibrium strategies, thereby extending the results of [14]. It should be noted that the equations for $V(k)$ are

given in implicit form, that is, they cannot be solved with respect to $V'(k)$, so that finding a C^2 solution requires special techniques (first a blow-up, and then the central manifold theorem).

We then consider the criterion:

$$(1 - \alpha) \delta \int_0^\infty e^{-\delta t} u(c(t), k(t)) dt + \alpha r \int_0^\infty e^{-rt} U(c(t), k(t)) dt \quad (10)$$

which belongs to the class (7) and we let $r \rightarrow 0$. In the limit, we get equilibrium strategies for the Chichilnisky problem (8), which we describe explicitly.

2 The Ramsey problem

This is the classical model for economic growth, originating with the seminal paper of Ramsey [31] in 1928, and developed by Cass [9], Koopmans [25] and many others (see [5] for a modern exposition). We are given a point $k_0 > 0$ and a concave continuous function f on $[0, \infty)$, which is C^∞ on $]0, \infty[$ and satisfies the Inada conditions:

$$\lim_{x \rightarrow 0} f'(x) = +\infty, \quad \lim_{x \rightarrow \infty} f'(x) \leq 0. \quad (11)$$

Definition 1 A capital-consumption path $(c(t), k(t))$, $t \geq 0$, is admissible if:

$$k(t) > 0 \text{ and } c(t) > 0 \text{ for all } t, \quad (12)$$

$$\frac{dk}{dt} = f(k) - c, \quad k(0) = k_0, \quad (13)$$

The set of all admissible paths starting from k_0 (i.e, such that $k(0) = k_0$) will be denoted by $\mathcal{A}(k_0)$.

We are given a number $\delta > 0$ and another concave, increasing function u on $]0, \infty)$, which is C^∞ on the interior, with $u''(c) > 0$ everywhere. We introduce the following criterion on $\mathcal{A}(k_0)$:

$$I(c, k) = \int_0^\infty u(c(t), k(t)) e^{-\delta t} dt, \quad (14)$$

and we consider the optimization problem:

$$\sup \{I(c, k) \mid (c, k) \in \mathcal{A}(k_0)\}. \quad (15)$$

The Euler-Lagrange equation is given by:

$$u''_{11} \frac{dc}{dt} = (\delta - f'(k)) u'_1 - u'_2 - (f(k) - c) u''_{12}. \quad (16)$$

Equation (16), together with equation (13), constitute a system of two first-order ODEs for the unknown functions $(c(t), k(t))$. In the particular case when $u = u(c)$ does not depend on k , the last equation simplifies to:

$$u''_{11} \frac{dc}{dt} = (\delta - f'(k)) u'_1$$

and the (16), (13) gives rise to a well-known phase diagram, with a hyperbolic stationary point (c_∞, k_∞) characterized by $f'(k_\infty) = \delta$ and $f(k_\infty) = c_\infty$. The optimal solution of the Ramsey

problem in that case then is the solution of (16), (13) which converges to (c_∞, k_∞) (see [5] for instance).

In the general case where $u(c, k)$ depends on k , the situation is not as simple, and to our knowledge has not been investigated. Stationary points (c_∞, k_∞) of the dynamics (if any) are given by:

$$c_\infty - f(k_\infty) = 0 \quad (17)$$

$$(\delta - f'(k_\infty)) u'_1(f(k_\infty), k_\infty) - u'_2(f(k_\infty), k_\infty) = 0. \quad (18)$$

To prove the existence of an optimal strategy, we do not use the Euler equation. We use the Hamilton-Jacobi-Bellman (HJB) equation instead. Introduce the optimal value as a function of the initial point:

$$V(k_0) := \sup \{I(c, k) \mid (c, k) \in \mathcal{A}(k_0)\}$$

If there is an optimal solution $(c(t), k(t))$, and it converges to (c_∞, k_∞) when $t \rightarrow \infty$, then, substituting in (14), we must have:

$$V(k_\infty) = \int_0^\infty e^{-\delta t} u(c_\infty, k_\infty) dt = \frac{1}{\delta} u(c_\infty, k_\infty) = \frac{1}{\delta} u(f(k_\infty), k_\infty) \quad (19)$$

Theorem 2 *If $V(k)$ is C^1 , it satisfies the HJB equation, namely:*

$$\delta V(k) = \max_c \{u(c, k) + (f(k) - c)V'(k)\}. \quad (20)$$

Conversely, suppose the HJB equation has a C^2 solution satisfying (19) for some (c_∞, k_∞) , and define a strategy $c = \sigma(k)$ by:

$$u'_1(\sigma(k), k) = V'(k) \quad (21)$$

Suppose moreover that the solution of:

$$\frac{dk}{dt} = f(k) - \sigma(k), \quad k(0) = k_0, \quad (22)$$

converges to k_∞ for all initial points k_0 . Then $\sigma(k)$ is an optimal solution of the generalized Ramsey problem (15)

This is the so-called verification theorem, which is classical (see [4]). We need V to be C^2 , so that σ is defined. Since $u''_1 > 0$, we can use the implicit function theorem on equation (21) to define σ . If V is C^2 , then σ is C^1 , and the initial-value problem (22) has a unique solution. Note also that everything is local: the functions V and σ are defined in some neighbourhood of k_∞ only, and the initial value k_0 is assumed to belong to that neighbourhood of k_∞ .

So, to prove the (local) existence of an optimal strategy $\sigma(k)$, we have to prove that the HJB equation has a C^2 solution V with $V(k_\infty) = \delta^{-1}u(f(k_\infty), k_\infty)$, and that the corresponding path $k(t)$ converges to k_∞ . Then the right-hand side of (22) converges to 0, so that $c(t) = \sigma(k(t))$ converges to $\sigma(k_\infty) = f(k_\infty)$. This is the content of the following two results

Theorem 3 Suppose there is some $k_\infty > 0$ satisfying (18) and:

$$u'_1 f'' + u''_{22} + f' \left(u''_{12} - \frac{u'_2}{u'_1} u''_{11} \right) - \frac{u'_2}{u'_1} u''_{12} < 0 \quad (23)$$

(all values to be taken at k_∞ and $c_\infty = f(k_\infty)$). Then there is an optimal strategy $c = \sigma(k)$ converging to k_∞

If $u = u(c)$ does not depend on k , then $u'_2 = 0$. Equation (18) becomes $f'(k_\infty) = \delta$, which defines k_∞ uniquely because of the Inada conditions (11) on f , and condition (23) becomes $u'_1(c_\infty) f''(k_\infty) < 0$, which is satisfied automatically. So there is an optimal strategy in that case, and one can even show that it is globally defined. For u depending on k , however, the situation is different.

Proof. We follow the method of [15]. By the inverse function theorem, the equation $u'_1(c, k) = p$ defines c as a C^1 function of p and k :

$$c = \varphi(p, k), \quad u'_1(\varphi(p, k), k) = p \quad (24)$$

We rewrite (20) as a Pfaff system:

$$dV = pdk, \quad (25)$$

$$p(f(k) - \varphi(p, k)) + u(\varphi(p, k), k) = \delta V, \quad (26)$$

and we seek a C^2 solution V satisfying:

$$V(k_\infty) = \frac{1}{\delta} u(f(k_\infty), k_\infty). \quad (27)$$

Differentiating (26) leads to:

$$\delta dV = (f(k) - \varphi(p, k))dp + pf'(k)dk + u'_2(\varphi(p, k), k)dk \quad (28)$$

Plugging (28) into (25), we get:

$$(f(k) - \varphi(p, k))dp = (\delta p - pf'(k) - u'_2(\varphi(p, k), k))dk. \quad (29)$$

Introducing an auxiliary variable t , we rewrite this as a system of two ODES for two functions $p(t)$ and $k(t)$:

$$\begin{aligned} \frac{dk}{dt} &= f(k) - \varphi(p, k) \\ \frac{dp}{dt} &= \delta p - pf'(k) - u'_2(\varphi(p, k), k) \end{aligned} \quad (30)$$

with the initial condition

$$(k(0), p(0)) = (k_\infty, p_\infty)$$

By (24), we must have

$$p_\infty = u'_1(\varphi(p_\infty, k_\infty), k_\infty) = u'_1(c_\infty, k_\infty) = u'_1(f(k_\infty), k_\infty)$$

Differentiating (24) with respect to p and k respectively, we derive the formulas:

$$\varphi'_1(p_\infty, k_\infty) = \frac{1}{u''_{11}(\varphi(p_\infty, k_\infty), k_\infty)} = \frac{1}{u''_{11}(f(k_\infty), k_\infty)}, \quad (31)$$

$$\varphi'_2(p_\infty, k_\infty) = -\frac{u''_{11}(\varphi(p_\infty, k_\infty), k_\infty)}{u''_{11}(\varphi(p_\infty, k_\infty), k_\infty)} = -\frac{u''_{12}(f(k_\infty), k_\infty)}{u''_{11}(f(k_\infty), k_\infty)}, \quad (32)$$

We can now linearizd (30) at the point (p_∞, k_∞) . We get:

$$\frac{d}{dt} \begin{pmatrix} k - k_\infty \\ p - p_\infty \end{pmatrix} = A_\infty \begin{pmatrix} k - k_\infty \\ p - p_\infty \end{pmatrix}.$$

where the constant matrix A_∞ is given by:

$$A_\infty := \begin{pmatrix} f' + \frac{u''_{12}}{u''_{11}}, & -\frac{1}{u''_{11}} \\ -u'_1 f'' - u''_{22} + \frac{(u''_{12})^2}{u''_{11}}, & \delta - f' - \frac{u''_{12}}{u''_{11}} \end{pmatrix}.$$

and all the values are to be taken at (k_∞, p_∞) . The characteristic polynomial is:

$$\lambda^2 - \delta\lambda - \left[\frac{u'_{1\infty} f''_\infty + u''_{22\infty}}{u''_{11\infty}} + f'_\infty \left(\frac{u''_{12\infty}}{u''_{11\infty}} - \frac{u'_{2\infty}}{u'_{1\infty}} \right) - \frac{u'_{2\infty}}{u'_{1\infty}} \frac{u''_{12\infty}}{u''_{11\infty}} \right] = 0$$

where we used $f'_\infty + \frac{u'_{2\infty}}{u'_{1\infty}} = \delta$ by (17). Because of assumption (23), it has two real roots with different signs, $\lambda_+ > 0$ and $\lambda_- < 0$. Thus (k_∞, p_∞) is a hyperbolic fixed point of (30), with a stable C^∞ -manifold \mathcal{S} which corresponds to λ_- and an unstable C^∞ -manifold \mathcal{U} which corresponds to λ_+ . Choose a smooth parametrization $(k_s(x), p_s(x))$ for the curve \mathcal{S} . The tangent at the fixed point is:

$$\frac{dp_s}{dk_s}(k_\infty) = u''_{11}(f'_\infty - \lambda_-) + u''_{12}$$

and plugging $k = k_s(x)$, $p = p_s(x)$ into equation (26), we get

$$V_s = \frac{p_s(x)(f(k_s(x)) - \varphi(p_s(x), k_s(x))) + u(\varphi(p_s(x), k_s(x)), k_s(x))}{\delta}.$$

Moreover, differentiating (26), and using (23) again, we find:

$$\frac{dV_s}{dk_s}(k_\infty) = u'_1(f(k_\infty), k_\infty) > 0,$$

It follows that the curve in parametric form $x \rightarrow (k(x), V(x))$ is in fact the graph of a function $V(k)$ which solves HJB and satisfies (27). By (25), we have

$$\frac{d^2 V_s}{dk_s^2}(k_\infty) = \frac{dp_s}{dk_s}(k_\infty) = u''_{11\infty}(f'_\infty - \lambda_-) + u''_{12\infty}. \quad (33)$$

Thus $V_s(k)$ is C^2 in k at k_∞ , and the C^2 property at other points near k_∞ follows from (30).

It remains to show that the strategy σ defined by $u'_1(\sigma(k), k) = V'(k)$ converges to $(f(k_\infty), k_\infty)$. We rewrite σ as:

$$\sigma(k) = \varphi(V'(k), k).$$

Linearizing the equation

$$\frac{dk}{dt} = f(k) - c = f(k) - \varphi(V'(k), k)$$

gives

$$\frac{d(k - k_\infty)}{dt} = \lambda_-(k - k_\infty)$$

and this concludes the proof. ■

Let us show how to deduce the Euler equation (16) from the HJB equation (20) and the optimal strategy (21). Setting $c(t) := \sigma(k(t))$, differentiating (20) with respect to k , and applying the envelope theorem, we get:

$$\delta V'(k) = u'_2(c, k) + f'(k) V'(k) + (f(k) - c) V''(k),$$

and hence, noting that $\frac{dk}{dt} = f(k) - c(t)$:

$$(\delta - f'(k)) V'(k) - u'_2(c, k) = (f(k) - c) \frac{d}{dk} V'(k) = \frac{d}{dt} V'(k).$$

Replacing $V'(k)$ by $u'_1(\sigma(k), k) = u'_1(c, k)$, we get:

$$(\delta - f'(k)) u'_1(c, k) - u'_2(c, k) = \frac{d}{dt} u'_1(c, k),$$

which is precisely the Euler equation.

3 Time-inconsistency.

3.1 Equilibrium strategies

We consider the intertemporal decision problem (as it seen at time $t = 0$)

$$J(c, k) = \int_0^\infty [h(t) u(c(t), k(t)) + H(t) U(c(t), k(t))] dt \quad (34)$$

under the dynamics described by (5) and (6). Here h and H are discount factors, i.e. C^∞ non-increasing functions on $[0, \infty)$, such that $h(0) = H(0) = 1$ and $h(\infty) = H(\infty) = 0$, while u and U are utility functions. They are assumed to be C^∞ on $]0, \infty)^2$, with $u'' < 0$ and $U'' < 0$ everywhere. We shall also assume that they decay exponentially, so that there is some $\rho > 0$ and some $T > 0$ such that $h(t) < e^{-\rho t}$ and $H(t) < e^{-\rho t}$ for $t \geq T$.

Because of time-inconsistency, the decision problem can no longer be seen as an optimization problem. There is no way for the decision-maker at time 0 to achieve what is, from her point of view, the first-best solution of the problem, and she must turn to a second-best policy: the best she

can do is to guess what her successor are planning to do, and then to plan her own consumption $c(0)$ accordingly. In other word, we will be looking for a subgame-perfect equilibrium of the leader-follower game played by successive generations.

The equilibrium policy was described in [14] for the case when the criterion (34) did not include the second term, and u did not depend on k . We will extend this analysis to the present situation, and then compare it with the approach in [22].

Definition 4 *A Markov strategy $c = \sigma(k)$ is convergent if there is a point k_∞ and a neighbourhood \mathcal{N} of k_∞ such that, for every $k_0 \in \mathcal{N}$ the solution $k(t)$ of (22) converges to k_∞ (and so $c(t) = \sigma(k(t))$ converges to $c_\infty = f(k_\infty)$). If this is the case, we shall say that (c_∞, k_∞) is supported by σ .*

Let us suppose that a convergent Markov strategy σ has been announced and is the public knowledge σ . The decision-maker at T has capital stock k_T . If all future decision-maker apply the strategy σ , the resulting future capital stock flow $k(t)$ obeys:

$$\frac{dk}{dt} = f(k) - \sigma(k), \quad k(T) = k_T, \quad T \leq t \quad (35)$$

Since every decision-maker faces the same problem (with different stock levels) it is enough to take $T = 0$. Suppose the decision-maker at time 0 holds power for $0 \leq t < \varepsilon$, and expects all later decision-makers to apply the strategy σ . He or she then explores whether it is in his or her interest to apply the strategy σ , that is, to play $c_0 = \sigma(k_0)$ for $0 \leq t < \varepsilon$. If he or she applies the constant control c for $0 \leq t \leq \varepsilon$.

Suppose the constant control c is use on $0 \leq t \leq \varepsilon$. The immediate utility flow during $[0, \varepsilon]$ is $[u(c, k_0) + U(c, k_0)]\varepsilon + o(\varepsilon)$ where $o(\varepsilon)$ is a higher order term of ε . At time ε , the resulting capital will be $k_0 + (f(k_0) - c)\varepsilon + o(\varepsilon)$. From then on, the strategy σ will be applied, which results in a capital stock k_c satisfying (we omit higher-order terms):

$$\frac{dk_c}{dt} = f(k_c) - \sigma(k_c), \quad k_c(\varepsilon) = k_0 + (f(k_0) - c)\varepsilon, \quad t \geq \varepsilon \quad (36)$$

The capital stock k_c can be written as

$$k_c(t) = k_0(t) + k_1(t)\varepsilon, \quad (37)$$

where $k_0(t)$ is the unperturbed solution, and $k_1(t)$ is given by the linearized equation:

$$\frac{dk_0}{dt} = f(k_0) - \sigma(k_0), \quad k_0(0) = k_0 \quad (38)$$

$$\frac{dk_1}{dt} = (f'(k_0) - \sigma'(k_0))k_1, \quad k_1(0) = \sigma(k_0) - c \quad (39)$$

Evaluating the integral (34) we get:

$$J(\varepsilon) = u(c, k_0)\varepsilon + \int_\varepsilon^\infty h(s)u(\sigma(k_0(t) + \varepsilon k_1(t)), k_0(t) + \varepsilon k_1(t))dt \\ + U(c, k_0)\varepsilon + \int_\varepsilon^\infty H(s)U(\sigma(k_0(t) + \varepsilon k_1(t)), k_0(t) + \varepsilon k_1(t))dt$$

Letting $\varepsilon \rightarrow 0$, so that commitment span of the decision-maker vanishes, we get:

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} (J(\varepsilon) - J(0)) = P(k_0, \sigma, c)$$

where

$$\begin{aligned}
P(k_0, \sigma, c) &= u(c, k_0) - u(\sigma(k_0), k_0) + (U(c, k_0) - U(\sigma(k_0), k_0)) \\
&+ \int_0^\infty h(t)u'_1(\sigma(k_0(t)), k_0(t))\sigma'(k_0(t))k_1(t)dt + \int_0^\infty h(t)u'_2(\sigma(k_0(t)), k_0(t))k_1(t)dt \\
&+ \int_0^\infty H(t)U'_1(\sigma(k_0(t)), k_0(t))\sigma'(k_0(t))k_1(t)dt + \int_0^\infty H(t)U'_2(\sigma(k_0(t)), k_0(t))k_1(t)dt,
\end{aligned}$$

Definition 5 A convergent Markov strategy σ is an equilibrium if we have:

$$\max_c P(k, \sigma, c) = P(k, \sigma, \sigma(k)), \quad \forall k.$$

3.2 The HJB approach

We now characterize the equilibrium strategy. We write $k_c(t) = \mathcal{K}(t; k_0, \sigma)$ where \mathcal{K} is the flow associated with the differential equation (38). We also define a function φ by:

$$\begin{aligned}
u'_1(\varphi(x, k), k) + U'_1(\varphi(x, k), k) &= x, \\
\varphi(u'_1(c, k) + U'_1(c, k), k) &= c
\end{aligned}$$

Since u''_1 and U''_1 are both negative, the function φ is well-defined by the implicit function theorem.

Theorem 6 Let σ be an equilibrium strategy. The function:

$$V(k_0) = \int_0^\infty h(t)u(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))dt + \int_0^\infty H(t)U(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))dt \quad (40)$$

satisfies the integral equation:

$$\begin{aligned}
V(k_0) &= \int_0^\infty h(t)u(\varphi \circ V'(\mathcal{K}(t; k_0, \varphi \circ V')), \mathcal{K}(t; k_0, \varphi \circ V'))dt \\
&+ \int_0^\infty H(t)U(\varphi \circ V'(\mathcal{K}(t; k_0, \varphi \circ V')), \mathcal{K}(t; k_0, \varphi \circ V'))dt
\end{aligned} \quad (\text{IE})$$

and the instantaneous optimality condition

$$u'_1(\sigma(k_0), k_0) + U'_1(\sigma(k_0), k_0) = V'(k_0), \quad \sigma(k_0) = \varphi(V'(k_0), k_0), \quad (41)$$

Conversely, suppose a function V is twice continuously differentiable, satisfies (IE), and the strategy $\sigma(k_0) := \varphi(V'(k_0), k_0)$ is convergent. Then σ is an equilibrium strategy.

For the sake of convenience, we have shortened $\sigma(k_0) := \varphi(V'(k_0), k_0)$ to $\sigma = \varphi \circ V'$.

Proof. Since the system is autonomous, we have:

$$\mathcal{K}(s; \mathcal{K}(t; k_0, \sigma), \sigma) = \mathcal{K}(s + t; k_0, \sigma). \quad (42)$$

Next, denote the fundamental solution of the linearized equation of (35) at k_0 by $\mathcal{R}(k_0; t)$ so that:

$$\begin{aligned}
k_1(t) &= \mathcal{R}(k_0; t)(\sigma(k_0) - c) \\
\frac{d\mathcal{R}}{dt} &= (f'(\mathcal{K}(t; k_0, \sigma)) - \sigma'(\mathcal{K}(t; k_0, \sigma)))\mathcal{R}(t), \quad \mathcal{R}(k_0; 0) = I,
\end{aligned}$$

\mathcal{R} and \mathcal{K} are related by:

$$\frac{\partial \mathcal{K}(t; k_0, \sigma)}{\partial k_0} = \mathcal{R}(k_0; t), \quad (43)$$

$$\frac{\partial \mathcal{K}(t; k_0, \sigma)}{\partial t} = f(\mathcal{K}(t; k_0, \sigma)) - \sigma(\mathcal{K}(t; k_0, \sigma)). \quad (44)$$

Let us now turn to the first part of the theorem. Differentiating (40) with respect to k_0 :

$$\begin{aligned} V'(k_0) &= \int_0^\infty h(t)u'_1(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))\sigma'(\mathcal{K}(t; k_0, \sigma))\mathcal{R}(k_0; t)dt \\ &\quad + \int_0^\infty h(t)u'_2(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))\mathcal{R}(k_0; t)dt \\ &\quad + \int_0^\infty H(t)U'_1(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))\sigma'(\mathcal{K}(t; k_0, \sigma))\mathcal{R}(k_0; t)dt \\ &\quad + \int_0^\infty H(t)U'_2(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))\mathcal{R}(k_0; t)dt. \end{aligned} \quad (45)$$

Substituting $k_0(t) = \mathcal{K}(t; k_0, \sigma)$ and $k_1(t)$ in the definition of P , we get:

$$\begin{aligned} P(k_0, \sigma, c) &= [u(c, k_0) + U(c, k_0)] - [u(\sigma(k_0), k_0) + U(\sigma(k_0), k_0)] \\ &\quad + \int_0^\infty h(t)u'_1(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))\sigma'(\mathcal{K}(t; k_0, \sigma))\mathcal{R}(k_0; t)(\sigma(k_0) - c)dt \\ &\quad + \int_0^\infty h(t)u'_2(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))\mathcal{R}(k_0; t)(\sigma(k_0) - c)dt \\ &\quad + \int_0^\infty H(t)U'_1(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))\sigma'(\mathcal{K}(t; k_0, \sigma))\mathcal{R}(k_0; t)(\sigma(k_0) - c)dt \\ &\quad + \int_0^\infty H(t)U'_2(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))\mathcal{R}(k_0; t)(\sigma(k_0) - c)dt. \end{aligned}$$

Since u and U are strictly concave and differentiable with respect to c , the necessary and sufficient condition to maximize $P(k_0, \sigma, c)$ with respect to c is that the derivative vanishes at $c = \sigma(k_0)$, that is:

$$\begin{aligned} u'_1(\sigma(k_0), k_0) + U'_1(\sigma(k_0), k_0) &= \int_0^\infty h(t)u'_1(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))\sigma'(\mathcal{K}(t; k_0, \sigma))\mathcal{R}(k_0; t)dt \\ &\quad + \int_0^\infty h(t)u'_2(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))\mathcal{R}(k_0; t)dt \\ &\quad + \int_0^\infty H(t)U'_1(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))\sigma'(\mathcal{K}(t; k_0, \sigma))\mathcal{R}(k_0; t)dt \\ &\quad + \int_0^\infty H(t)U'_2(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))\mathcal{R}(k_0; t)dt, \end{aligned}$$

The right-hand side is precisely $V'(k_0)$, as we wanted. Therefore, the equilibrium strategy satisfies

$$u'_1(\sigma(k_0), k_0) + U'_1(\sigma(k_0), k_0) = V'(k_0)$$

and we have $\sigma(k_0) = \varphi(V'(k_0), k_0)$. Substituting back into equation (40), we get the functional equation (IE). This proves the first part of the theorem (necessity). We refer to [14] for the second part (sufficiency). ■

The following theorem gives an alternative characterization, the differential equation, which resembles the usual HJB equation from the calculus of variation.

Theorem 7 *Let V be a C^2 function such that the strategy $\sigma = \varphi \circ V'$ converges to \bar{k} . Then V satisfies the integral equation (IE) if and only if it satisfies the following integro-differential equation*

$$\begin{aligned} & u(\varphi \circ V'(k_0), k_0) + U(\varphi \circ V'(k_0), k_0) + V'(k_0)(f(k_0) - \varphi(V'(k_0), k_0)) \\ &= - \int_0^\infty h'(t)u(\varphi \circ V'(\mathcal{K}(t; k_0, \varphi \circ V')), \mathcal{K}(t; k_0, \varphi \circ V'))dt \\ & \quad - \int_0^\infty H'(t)U(\varphi \circ V'(\mathcal{K}(t; k_0, \varphi \circ V')), \mathcal{K}(t; k_0, \varphi \circ V'))dt \end{aligned} \quad (\text{DE})$$

together with the boundary condition

$$V(\bar{k}) = u(f(\bar{k}), \bar{k}) \int_0^\infty h(t)dt + U(f(\bar{k}), \bar{k}) \int_0^\infty H(t)dt. \quad (\text{BC})$$

Proof. Introduce the function ϕ defined by

$$\phi(k_0) = V(k_0) - \int_0^\infty h(t)u(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))dt - \int_0^\infty H(t)U(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))dt,$$

where $\sigma(k_0) = \varphi(V'(k_0), k_0)$. Consider the value $\psi(t, k_0)$ of ϕ along the trajectory $t \rightarrow \mathcal{K}(t; k_0, \sigma)$ originating from k_0 at time 0, that is

$$\begin{aligned} \psi(t, k_0) &= \phi(\mathcal{K}(t; k_0, \sigma)) \\ &= V(\mathcal{K}(t; k_0, \sigma)) - \int_t^\infty h(s-t)u(\sigma(\mathcal{K}(s; k_0, \sigma)), \mathcal{K}(s; k_0, \sigma))ds \\ & \quad - \int_t^\infty H(s-t)U(\sigma(\mathcal{K}(s; k_0, \sigma)), \mathcal{K}(s; k_0, \sigma))ds, \end{aligned}$$

Then the derivative of ψ with respect to t is given by:

$$\begin{aligned} \frac{\partial \psi(t, k_0)}{\partial t} &= V'(\mathcal{K}_t)[f(\mathcal{K}_t) - \varphi \circ V'(\mathcal{K}_t)] + u(\varphi \circ V'(\mathcal{K}_t), \mathcal{K}_t) + U(\varphi \circ V'(\mathcal{K}_t), \mathcal{K}_t) \\ & \quad + \int_0^\infty h'(s)u(\varphi \circ V'(\mathcal{K}(s; \mathcal{K}_t, \varphi \circ V')), \mathcal{K}(s; \mathcal{K}_t, \varphi \circ V'))ds \\ & \quad + \int_0^\infty H'(s)U(\varphi \circ V'(\mathcal{K}(s; \mathcal{K}_t, \varphi \circ V')), \mathcal{K}(s; \mathcal{K}_t, \varphi \circ V'))ds, \end{aligned}$$

where we have denoted $\mathcal{K}(t; k_0, \varphi \circ V')$ by \mathcal{K}_t for convenience. If (DE) holds, then the right hand side is identically zero along the trajectory, so that $\psi(t, k_0)$ does not depend on t , thus $\psi(s, k_0) = \psi(t, k_0)$ for all $s, t \geq 0$. Letting $t \rightarrow \infty$ in the definition of ψ , we get

$$\begin{aligned} \psi(s, k_0) &= \lim_{t \rightarrow \infty} \psi(t, k_0) \\ &= V(\bar{k}) - \int_0^\infty h(s)u(\sigma(\bar{k}), \bar{k})ds - \int_0^\infty H(s)U(\sigma(\bar{k}), \bar{k})ds, \end{aligned} \quad (46)$$

and hence, if (BC) holds, then $\psi = \phi \equiv 0$ and so equation (IE) holds. Conversely, if $V(k)$ satisfies equation (IE), then the same lines of reasoning shows that equation (DE) and the boundary condition (BC) are satisfied. ■

3.3 The Euler equations

To obtain the Euler-Lagrange-like equation, we differentiate the both side of (DE) with respect to k_0 . We get:

$$\begin{aligned}
& - \int_0^\infty h'(t)u'_1(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))\sigma'(\mathcal{K}(t; k_0, \sigma))\frac{\partial\mathcal{K}(t; k_0, \sigma)}{\partial k_0}dt \\
& - \int_0^\infty h'(t)u'_2(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))\frac{\partial\mathcal{K}(t; k_0, \sigma)}{\partial k_0}dt \\
& - \int_0^\infty H'(t)U'_1(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))\sigma'(\mathcal{K}(t; k_0, \sigma))\frac{\partial\mathcal{K}(t; k_0, \sigma)}{\partial k_0}dt \\
& - \int_0^\infty H'(t)U'_2(\sigma(\mathcal{K}(t; k_0, \sigma)), \mathcal{K}(t; k_0, \sigma))\frac{\partial\mathcal{K}(t; k_0, \sigma)}{\partial k_0}dt \\
& = [u'_1(\sigma(k_0), k_0) + U'_1(\sigma(k_0), k_0)]\sigma'(k_0) + u'_2(\sigma(k_0), k_0) + U'_2(\sigma(k_0), k_0) \\
& \quad + V'(k_0)f'(k_0) - V'(k_0)\sigma'(k_0) + V''(k_0)(f(k_0) - \sigma(k_0)).
\end{aligned}$$

Plugging $k_0 = k(t)$, $\sigma(k(t)) = c(t)$ and using (41) to cancel the first and the fifth terms, together with (13) and (43), we have

$$\begin{aligned}
& - \int_0^\infty h'(s)u'_1(\sigma(\mathcal{K}(s; k(t), \sigma)), \mathcal{K}(s; k(t), \sigma))\sigma'(\mathcal{K}(s; k(t), \sigma))\mathcal{R}(k(t), s)ds \\
& - \int_0^\infty h'(s)u'_2(\sigma(\mathcal{K}(s; k(t), \sigma)), \mathcal{K}(s; k(t), \sigma))\mathcal{R}(k(t), s)ds \\
& - \int_0^\infty H'(s)U'_1(\sigma(\mathcal{K}(s; k(t), \sigma)), \mathcal{K}(s; k(t), \sigma))\sigma'(\mathcal{K}(s; k(t), \sigma))\mathcal{R}(k(t), s)ds \\
& - \int_0^\infty H'(s)U'_2(\sigma(\mathcal{K}(s; k(t), \sigma)), \mathcal{K}(s; k(t), \sigma))\mathcal{R}(k(t), s)ds \\
& = u'_2(c(t), k(t)) + U'_2(c(t), k(t)) + V'(k(t))f'(k(t)) + V''(k(t))(f(k(t)) - c(t)) \\
& = u'_2(c(t), k(t)) + U'_2(c(t), k(t)) + [u'_1(c(t), k(t)) + U'_1(c(t), k(t))]f'(k(t)) \\
& \quad + \frac{d}{dt}[u'_1(c(t), k(t)) + U'_1(c(t), k(t))],
\end{aligned}$$

where to get the last two terms in the right hand side, we have used (41) again.

We have $\mathcal{K}(s; k(t), \sigma) = k(s+t)$, and $\mathcal{R}(k(t); s) = \mathcal{R}(k_0; s+t)$

$$\begin{aligned}
\sigma'(k(t)) &= \frac{1}{f(k(t)) - c(t)} \frac{dc}{dt} \\
\mathcal{R}(k_0; s+t) &= \exp(f'(k(t)) - \frac{1}{f(k(t)) - c(t)} \frac{dc}{dt}),
\end{aligned} \tag{47}$$

Writing this into the preceding equations, we finally get:

$$- \int_t^\infty h'(s-t) [u'_1(c(s), k(s))\gamma(s) + u'_2(c(s), k(s))] e^{f'(k(s))-\gamma(s)} ds \quad (48)$$

$$- \int_t^\infty H'(s-t) [U'_1(c(s), k(s))\gamma(s) + U'_2(c(s), k(s))] e^{f'(k(s))-\gamma(s)} ds \quad (49)$$

$$= u'_2(c(t), k(t)) + U'_2(c(t), k(t)) + [u'_1(c(t), k(t)) + U'_1(c(t), k(t))] f'(k(t)) \quad (50)$$

$$+ \frac{d}{dt} [u'_1(c(t), k(t)) + U'_1(c(t), k(t))] \quad (51)$$

with:

$$\gamma(s) := \frac{1}{f(k(s)) - c(s)} \frac{dc}{ds} \quad (52)$$

which is the Euler-Lagrange-like equation for the time-inconsistent case.

3.4 The control theory approach

Karp [22], [23] has developed a different method to deal with time-inconsistency. In this section, we connect his results with ours.

Defining $V(k_0)$ as above, we must have:

$$V(k_0) = \max_c \left\{ u(c, k_0) + U(c, k_0) \varepsilon + \int_\varepsilon^\infty [h(t)u(\sigma(k_c(t)), k_c(t)) + H(t)U(\sigma(k_c(t)), k_c(t))] dt \right\} \quad (53)$$

On the other hand, we have

$$V(k_c(\varepsilon)) = \int_\varepsilon^\infty [h(s-\varepsilon)u(\sigma(k_c(s)), k_c(s)) + H(s-\varepsilon)U(\sigma(k_c(s)), k_c(s))] ds,$$

Substituting:

$$V(k_0) = \max_c \left\{ \begin{array}{l} [u(c, k_0) + U(c, k_0)] \varepsilon + V(k_c(\varepsilon)) \\ + \int_\varepsilon^\infty [h(t) - h(t-\varepsilon)] u(\sigma(k_c(t)), k_c(t)) dt \\ + \int_\varepsilon^\infty [H(t) - H(t-\varepsilon)] U(\sigma(k_c(t)), k_c(t)) dt \end{array} \right\}$$

Letting $\varepsilon \rightarrow 0$, we have

$$\begin{aligned} & - \int_0^\infty h'(t)u(\sigma(k(t)), k(t)) dt - \int_0^\infty H'(t)U(\sigma(k(t)), k(t)) dt \\ & = \max_c \{ u(c, k_0) + U(c, k_0) + V'(k_0)(f(k_0) - c) \}. \end{aligned} \quad (54)$$

This equation was first obtained by Karp [22], [23]. This is the HJB equation for a certain control problem, which he terms the auxiliary control problem. We now show that this approach is equivalent to the preceding one. approach by Ekeland-Lazrak. In the right hand side of (54), the maximum is attained at:

$$c = \arg \max_c \{ u(c, k_0) + U(c, k_0) + V'(k_0)(f(k_0) - c) \} = \varphi(V'(k_0), k_0),$$

thus we have

$$\begin{aligned} & - \int_0^\infty h'(t)u(\sigma(k(t)), k(t))dt - \int_0^\infty H'(t)U(\sigma(k(t)), k(t))dt \\ & = u(\varphi(V'(k_0), k_0), k_0) + U(\varphi(V'(k_0), k_0), k_0) + V'(k_0)(f(k_0) - \varphi(V'(k_0), k_0))). \end{aligned}$$

which is exactly the same as (DE)

4 The biexponential case

4.1 The equations

In this section, we consider the biexponential criterion

$$\lambda \int_0^\infty e^{-\delta_1 s} u(c(s), k(s)) ds + (1 - \lambda) \int_0^\infty e^{-\delta_2 s} U(c(s), k(s)) ds. \quad (55)$$

Without loss of generality, we assume that:

$$\delta_1 > \delta_2$$

If $\lambda = 0$ or 1 , then this is just the Ramsey criterion (14). Thus we are interested in the case $0 < \lambda < 1$. Dividing by λ , we find that (55) is a special case of (34), where $h(t) = e^{-\delta_1 t}$, $H(t) = e^{-\delta_2 t}$ and U is replaced by $\frac{1-\lambda}{\lambda}U$. So all the results of the preceding section hold.

4.1.1 The HJB-type equations

Given an equilibrium strategy σ_λ , introduce the two functions:

$$V_\lambda(k) := \int_0^\infty e^{-\delta_1 t} u(\sigma_\lambda(k(t)), k(t)) dt + \frac{1-\lambda}{\lambda} \int_0^\infty e^{-\delta_2 t} U(\sigma_\lambda(k(t)), k(t)) dt \quad (56)$$

$$W_\lambda(k) := \int_0^\infty e^{-\delta_1 t} u(\sigma_\lambda(k(t)), k(t)) dt - \frac{1-\lambda}{\lambda} \int_0^\infty e^{-\delta_2 t} U(\sigma_\lambda(k(t)), k(t)) dt \quad (57)$$

In Proposition 8, we prove that the HJB-type equation (DE) reduces to a system of two ODEs for V_λ and W_λ :

$$(f - \varphi_\lambda(V'_\lambda)) V'_\lambda + u(V'_\lambda) + \frac{1-\lambda}{\lambda} U(V'_\lambda) = \frac{\delta_1 + \delta_2}{2} V_\lambda + \frac{\delta_1 - \delta_2}{2} W_\lambda, \quad (58)$$

$$(f - \varphi_\lambda(V'_\lambda)) W'_\lambda + u(V'_\lambda) - \frac{1-\lambda}{\lambda} U(V'_\lambda) = \frac{\delta_1 - \delta_2}{2} V_\lambda + \frac{\delta_1 + \delta_2}{2} W_\lambda \quad (59)$$

where $\varphi_\lambda(V'_\lambda)$ and $u(V'_\lambda)$ denote the functions $k \rightarrow \varphi_\lambda(V'_\lambda(k), k)$ and $k \rightarrow u(\varphi_\lambda(V'), k)$. Recall that φ_λ is defined by the equivalent equations:

$$u'_1(\varphi_\lambda(x, k), k) + \frac{1-\lambda}{\lambda} U'_1(\varphi_\lambda(x, k), k) = x, \quad (60)$$

$$\varphi_\lambda(u'_1(c, k) + \frac{1-\lambda}{\lambda} U'_1(c, k), k) = c \quad (61)$$

Similarly, the boundary condition (BC) becomes:

$$V_\lambda(k_\infty) = \frac{1}{\delta_1} u(f(k_\infty), k_\infty) + \frac{1-\lambda}{\lambda} \frac{1}{\delta_2} U(f(k_\infty), k_\infty), \quad (62)$$

$$W_\lambda(k_\infty) = \frac{1}{\delta_1} u(f(k_\infty), k_\infty) - \frac{1-\lambda}{\lambda} \frac{1}{\delta_2} U(f(k_\infty), k_\infty). \quad (63)$$

If $\lambda = 1$, we find the usual HJB equation (20) for V with the boundary condition (19).

Proposition 8 *Suppose that σ_λ is an equilibrium strategy such that $k(t)$ converges to k_∞ . Then V_λ and W_λ defined by (56) and (57) satisfy the equations (58) and (59) with the boundary conditions (62) and (63). Conversely, suppose there is a point k_∞ , a C^2 function V_λ and a C^1 function W_λ , both defined on some open neighbourhood of k_∞ , satisfying the equations (58) and (59) with the boundary conditions (62) and (63); suppose moreover that strategy $\sigma_\lambda(k) := \varphi_\lambda(V'_\lambda(k), k)$ converges to $(f(k_\infty), k_\infty)$. Then σ_λ is an equilibrium strategy.*

Let us draw the reader's attention to the fact that V_λ must be C^2 while W_λ needs only be C^1 .

Proof. Let us simplify the notation. Write σ instead of σ_λ and set

$$a = \frac{\delta_1 + \delta_2}{2}, \quad b = \frac{\delta_1 - \delta_2}{2}$$

Arguing as in Theorem 7, we obtain (58) and (59) by differentiating (56) and (57). The boundary conditions (62) and (63) follow from setting $k(t) = k_\infty$ and $c(t) = \sigma(k_\infty)$ in (58) and (59).

Conversely suppose v_1 and w_1 satisfy (58), (59) and (62), (63), and suppose the strategy $\sigma_1 = \varphi_\lambda \circ v'_1$ converges to k . Consider the following functions

$$\begin{aligned} v_2(k_0) &= \int_0^\infty e^{-\delta_1 t} u(\sigma_1(\mathcal{K}(t; k_0, \sigma_1)), \mathcal{K}(t; k_0, \sigma_1)) dt \\ &\quad + \frac{1-\lambda}{\lambda} \int_0^\infty e^{-\delta_2 t} U(\sigma_1(\mathcal{K}(t; k_0, \sigma_1)), \mathcal{K}(t; k_0, \sigma_1)) dt, \\ w_2(k_0) &= \int_0^\infty e^{-\delta_1 t} u(\sigma_1(\mathcal{K}(t; k_0, \sigma_1)), \mathcal{K}(t; k_0, \sigma_1)) dt \\ &\quad - \frac{1-\lambda}{\lambda} \int_0^\infty e^{-\delta_2 t} U(\sigma_1(\mathcal{K}(t; k_0, \sigma_1)), \mathcal{K}(t; k_0, \sigma_1)) dt. \end{aligned}$$

Arguing as in Theorem 7, we find that v_2 and w_2 also satisfy (58), (59) and (62), (63). Setting $v_3 = v_1 - v_2, w_3 = w_1 - w_2$, then we get

$$\begin{aligned} (f(k) - \sigma(k)) v'_3(k) + u(\sigma(k), k) + \frac{1-\lambda}{\lambda} U(\sigma(k), k) &= a v_3(k) + b w_3(k) \\ (f(k) - \sigma(k)) w'_3(k) + u(\sigma(k), k) + \frac{1-\lambda}{\lambda} U(\sigma(k), k) &= b v_3(k) + a w_3(k) \end{aligned} \quad (64)$$

with the boundary conditions

$$\begin{aligned} v_3(k_\infty) &= 0 \\ w_3(k_\infty) &= 0 \end{aligned} \quad (65)$$

Obviously, $v_3 = w_3 = 0$ is a solution. Lemma 9 below shows that We need to show that it is the only one, so that $v_1 = v_2$ and $w_1 = w_2$. This is the desired result. ■

Lemma 9 *If (v_3, w_3) is a pair of continuous functions on a neighborhood k_∞ , continuously differentiable for $k \neq k_\infty$, and which solve (64) with boundary conditions (65), then $v_3 = w_3 = 0$.*

Proof. Set $f(k) - \sigma_1(k) = \xi(k)$, then $\xi(k) \rightarrow 0$ as $k \rightarrow k_\infty$. The system (64) can be rewritten as:

$$\begin{pmatrix} \xi v_3' \\ \xi w_3' \end{pmatrix} = \begin{pmatrix} a & b \\ b & a \end{pmatrix} \begin{pmatrix} v_3 \\ w_3 \end{pmatrix}.$$

Setting:

$$\begin{pmatrix} V \\ W \end{pmatrix} = \begin{pmatrix} \sqrt{\frac{1}{2}} & \sqrt{\frac{1}{2}} \\ -\sqrt{\frac{1}{2}} & \sqrt{\frac{1}{2}} \end{pmatrix} \begin{pmatrix} v_3 \\ w_3 \end{pmatrix},$$

we have:

$$\xi \begin{pmatrix} V' \\ W' \end{pmatrix} = \begin{pmatrix} \delta_1 & 0 \\ 0 & \delta_2 \end{pmatrix} \begin{pmatrix} V \\ W \end{pmatrix}.$$

The first equation yields $V(k) = V(k_0) \exp \int_{k_0}^k \frac{\delta_1}{\xi(u)} du$. Without loss of generality, we can assume that $k_0 < k_\infty$.

Let $S = \{k \mid \xi(k) = 0, k_0 \leq k \leq k_\infty\}$, where k_0 is the initial stock. We consider the following two cases:

First case: $S = \{k_\infty\}$. Then $\xi(k) > 0$ for $k \in [k_0, k_\infty)$. So $\frac{dk}{dt} = \xi(k) > 0$, and:

$$0 = \sqrt{\frac{1}{2}}(v_3(k_\infty) + w_3(k_\infty)) = V(k_\infty) = V(k_0) \exp \int_{k_0}^{k_\infty} \frac{\delta_1}{\xi(u)} du$$

It follows that $V(k) = 0$ for all k .

Second case: $S \setminus \{k_\infty\} \neq \emptyset$. Then S is a closed set and $V(k) = \frac{\xi(k)V'(k)}{\delta_1} = 0$ in S . The complement of S is a countable union of disjoint intervals, and ξ vanishes at the endpoints of each interval. Arguing as above it follows that $V(k)$ vanishes for $k_0 \leq k \leq k_\infty$.

The same argument holds for $W(k)$. This concludes the proof. ■

4.1.2 The Euler-type equations

The Euler-type equations (48) to (52) can be reduced to a non-autonomous system of four first-order ODEs. It is better to work directly on (58) and (59) to get an autonomous system. Proceeding as in the end of Section 2, we differentiate (58) and (59) with respect to k , and notice that

$$(f - \sigma)V_\lambda'' = \frac{dk}{dt} \frac{V_\lambda'}{dk} = \frac{d}{dt} V_\lambda',$$

and then get:

$$\begin{aligned} \frac{\delta_1 + \delta_2}{2} V_\lambda' + \frac{\delta_1 - \delta_2}{2} W_\lambda' &= \frac{d}{dt} V_\lambda' + (f' - \sigma') V_\lambda' + (u_1' + \frac{1-\lambda}{\lambda} U_1') \sigma' + u_2' + \frac{1-\lambda}{\lambda} U_2', \\ \frac{\delta_1 - \delta_2}{2} V_\lambda' + \frac{\delta_1 + \delta_2}{2} W_\lambda' &= \frac{d}{dt} W_\lambda' + (f' - \sigma') W_\lambda' + (u_1' - \frac{1-\lambda}{\lambda} U_1') \sigma' + u_2' - \frac{1-\lambda}{\lambda} U_2', \end{aligned}$$

where $c = \varphi(V'_\lambda(k), k)$ and σ' is given by (47). Using formula (60), and setting $w(t) := W'_\lambda(k(t))$, this becomes:

$$\begin{aligned} \frac{\delta_1 + \delta_2}{2} \left(u'_1 + \frac{1-\lambda}{\lambda} U'_1 \right) + \frac{\delta_1 - \delta_2}{2} w &= \frac{d}{dt} \left(u'_1 + \frac{1-\lambda}{\lambda} U'_1 \right) + (f' - \sigma') \left(u'_1 + \frac{1-\lambda}{\lambda} U'_1 \right) \\ &\quad + \left(u'_1 + \frac{1-\lambda}{\lambda} U'_1 \right) \sigma' + u'_2 + \frac{1-\lambda}{\lambda} U'_2, \\ \frac{\delta_1 - \delta_2}{2} \left(u'_1 + \frac{1-\lambda}{\lambda} U'_1 \right) + \frac{\delta_1 + \delta_2}{2} w &= \frac{dw}{dt} + (f' - \sigma') w \\ &\quad + \left(u'_1 - \frac{1-\lambda}{\lambda} U'_1 \right) \sigma' + u'_2 - \frac{1-\lambda}{\lambda} U'_2, \end{aligned}$$

and hence

$$\left(\frac{\delta_1 + \delta_2}{2} - f' \right) (\lambda u'_1 + (1-\lambda) U'_1) + \frac{\delta_1 - \delta_2}{2} \lambda w = \frac{d}{dt} (\lambda u'_1 + (1-\lambda) U'_1) + \lambda u'_2 + (1-\lambda) U'_2, \quad (66)$$

$$\begin{aligned} \frac{\delta_1 - \delta_2}{2} (\lambda u'_1 + (1-\lambda) U'_1) + \left(\frac{\delta_1 + \delta_2}{2} - f' \right) \lambda w &= \lambda \frac{dw}{dt} + (-\lambda w + \lambda u'_1 - (1-\lambda) U'_1) \sigma' \\ &\quad + \lambda u'_2 - (1-\lambda) U'_2, \end{aligned} \quad (67)$$

to which we should add:

$$\frac{dk}{dt} = f(k) - c. \quad (68)$$

The system (66) to (68) is a system of three first-order ODEs for the unknown functions $k(t)$, $c(t)$ and $w(t)$. Note that for $\lambda = 0$, (66) and (67) reduce to two copies of the usual Euler-Lagrange equation. For $\lambda = 1$, taking $w(t) = u'_1(c(t), k(t))$ gives us again two copies of the same equation.

4.1.3 The control theory approach

Equation (54) is the HJB equation for the following:

$$\max_{c(\cdot)} \int_0^\infty e^{-rt} \left[u(c(t), k(t)) + \frac{1-\lambda}{\lambda} U(c(t), k(t)) - K(k(t)) \right] dt \quad (69)$$

$$\frac{dk}{dt} = f(k) - c(t), \quad k(0) = k_0 \quad (70)$$

where we seek a feedback $\sigma(k)$ such that:

$$K(k_0) = \left\{ (\delta - r) \int_0^\infty e^{-\delta t} u(\sigma(k(t)), k(t)) dt \mid \frac{dk}{dt} = f(k) - \sigma(k), k(0) = k_0 \right\} \quad (71)$$

Solving (69), (70) under the constraint (71) is a fixed-point problem for the feedback $c = \sigma(k)$.

4.2 Solving the boundary-value problem

Define a function $\bar{g}_\lambda(k)$ by:

$$\bar{g}_\lambda(k) := \frac{\lambda \delta_1 u'_1(f(k), k) + (1-\lambda) \delta_2 U'_1(f(k), k)}{\lambda u'_1(f(k), k) + (1-\lambda) U'_1(f(k), k)} - \frac{\lambda u'_2(f(k), k) + (1-\lambda) U'_2(f(k), k)}{\lambda u'_1(f(k), k) + (1-\lambda) U'_1(f(k), k)} \quad (72)$$

Theorem 10 Assume $\bar{g}_\lambda(k) \neq 0$. If there is some k_∞ such that

$$f'(k_\infty) \neq \bar{g}_\lambda(k_\infty), \quad (73)$$

then the equations (58) and (59) with the boundary conditions (62) and (63) have a solution (V, W) near the point k_∞ with V of class C^2 and W of class C^1 .

Proof. We adapt the argument in [14] to the present situation. We note that the boundary-value problem (58), (59), (62), (63) cannot be reduced to a standard initial-value problem for the pair (V_λ, W_λ) . To see that, rewrite equation (58) as follows:

$$(f(k) - \varphi(V'_\lambda, k)) V'_\lambda + u(\varphi(V'_\lambda, k), k) + \frac{1-\lambda}{\lambda} U(\varphi(V'_\lambda, k), k) = aV_\lambda + bW_\lambda.$$

Since the function $c \rightarrow u(c, k) + \frac{1-\lambda}{\lambda} U(c, k)$ is concave, the function:

$$c \rightarrow u(c, k) + \frac{1-\lambda}{\lambda} U(c, k) - cx$$

attains its maximum at the point $c = \varphi_\lambda(x, k)$ defined by (61). We set:

$$u_\lambda^*(x, k) = \max_c \left\{ u(c, k) + \frac{1-\lambda}{\lambda} U(c, k) - cx \right\} = u(\varphi_\lambda(x, k), k) + \frac{1-\lambda}{\lambda} U(\varphi_\lambda(x, k), k) - \varphi_\lambda(x, k)x \quad (74)$$

The function $x \rightarrow u_\lambda^*(x, k)$ is convex, and the equation (58) becomes:

$$f(k) V'_\lambda + u_\lambda^*(V'_\lambda, k) = aV_\lambda + bW_\lambda. \quad (75)$$

This is an equation for V'_λ . From the basic duality results in convex analysis (see for instance [17]), we find that:

$$\min_y \{f(k)y + u_\lambda^*(y, k)\} = u(f(k), k) + \frac{1-\lambda}{\lambda} U(f(k), k).$$

Note that $f(k)y + u_\lambda^*(y, k)$ is convex in y with minimal value $u(f(k), k) + \frac{1-\lambda}{\lambda} U(f(k), k)$. Then (75), considered as an equation for V'_λ , has two solutions if:

$$aV_\lambda(k) + bW_\lambda(k) > u(f(k), k) + \frac{1-\lambda}{\lambda} U(f(k), k),$$

and no solutions if the inverse inequality holds. If we have equality:

$$aV_\lambda(k) + bW_\lambda(k) = u(f(k), k) + \frac{1-\lambda}{\lambda} U(f(k), k)$$

then equation (75) has precisely one solution, namely:

$$V'_\lambda = u'_1(f(k), k) + \frac{1-\lambda}{\lambda} U'_1(f(k), k)$$

At the point k_∞ , with the values (62), (63), we find that:

$$aV_\lambda(k_\infty) + bW_\lambda(k_\infty) = u(f(k_\infty), k_\infty) + \frac{1-\lambda}{\lambda} U(f(k_\infty), k_\infty),$$

so that we are exactly on the boundary case. Equation (75) has precisely one solution, by (84) below, which satisfies

$$V'_\lambda(k_\infty) = u'_1(f(k_\infty), k_\infty) + \frac{1-\lambda}{\lambda} U'_1(f(k_\infty), k_\infty),$$

but it is degenerate:

$$\frac{\partial}{\partial y} [f(k_\infty)y + u^*_\lambda(y, k_\infty)]|_{y=V'_\lambda(k_\infty)} = 0,$$

so that equation (75) cannot be written in the form $V'_\lambda = \psi(k, V_\lambda)$. It is an implicit differential equation, and special techniques are needed to solve it. The difficulty is further increased by the fact that we need only V (but not W) to be C^2 .

The proof of Theorem 10 proceeds in several steps.

Step 1: Changing the unknown functions from $(V_\lambda(k), W_\lambda(k))$ to $(V_\lambda(k), \mu(k))$

Using (74), we find that the equation (58) is equivalent to

$$F(V'(k), k) = \mu(k)$$

with:

$$\begin{aligned} F(x, k) &:= u^*_\lambda(x, k) + xf(k) - u(f(k), k) - \frac{1-\lambda}{\lambda} U(f(k), k) \\ \mu(k) &:= aV_\lambda(k) + bW_\lambda(k) - u(f(k), k) - \frac{1-\lambda}{\lambda} U(f(k), k) \end{aligned}$$

Note that $x \rightarrow F(x, k)$ is a linear perturbation of the convex function $x \rightarrow u^*_\lambda(x, k)$. To shorten the notations, let us set:

$$y^*(k) := u'_1(f(k), k) + \frac{1-\lambda}{\lambda} U'_1(f(k), k) \quad (76)$$

As we pointed out, the equation

$$F(x + y^*(k), k) = \mu$$

in the variable x has two solutions $x_-(k, \mu) < 0 < x_+(k, \mu)$ for $\mu(k) > 0$, none for $\mu(k) < 0$ and a single solution $x = 0$ for $\mu(k) = 0$.

From the equation (59), we have

$$W'_\lambda = V'_\lambda \frac{bV_\lambda + aW_\lambda - u(\sigma_\lambda(k), k) + \frac{1-\lambda}{\lambda} U(\sigma_\lambda(k), k)}{aV_\lambda + bW_\lambda - u(\sigma_\lambda(k), k) - \frac{1-\lambda}{\lambda} U(\sigma_\lambda(k), k)}$$

Differentiating $\mu(k)$ with respect to k yields

$$\begin{aligned} \frac{d\mu}{dk} &= V'_\lambda \frac{(a^2 + b^2)V_\lambda + 2abW_\lambda - \delta_1 u(\sigma_\lambda(k), k) - \frac{(1-\lambda)\delta_2}{\lambda} U(\sigma_\lambda(k), k)}{aV_\lambda + bW_\lambda - u(\sigma_\lambda(k), k) - \frac{1-\lambda}{\lambda} U(\sigma_\lambda(k), k)} \\ &\quad - y^* f' - u'_2(f(k), k) - \frac{1-\lambda}{\lambda} U'_2(f(k), k) \end{aligned}$$

We now take $(V_\lambda(k), \mu(k))$ as our new unknown functions. They satisfy the equations:

$$\begin{aligned}\frac{dV}{dk} &= y^*(k) + x(k, \mu(k)), \\ \frac{d\mu}{dk} &= (y^* + x(k, \mu(k))) \frac{(a^2 + b^2)V_\lambda + 2abW_\lambda - \delta_1 u(\sigma_\lambda(k), k) - \frac{(1-\lambda)\delta_2}{\lambda} U(\sigma_\lambda(k), k)}{aV_\lambda + bW_\lambda - u(\sigma_\lambda(k), k) - \frac{1-\lambda}{\lambda} U(\sigma_\lambda(k), k)} \\ &\quad - y^* f' - u'_2(f(k), k) - \frac{1-\lambda}{\lambda} U'_2(f(k), k).\end{aligned}$$

In fact, according to which determination $(x(k, \mu(k)))$ is chosen, $x_+(k, \mu(k))$ or $x_-(k, \mu(k))$, these equations define two distinct dynamical systems on the region $\mu > 0$.

Step 3: Taking x instead of k as the independent variable.

To get rid of the indetermination, we pick x instead of k as the independent variable. We shorten our notation by setting $x(k) = x(\mu(k), k)$. We get:

$$\frac{dk}{dx} = \frac{f(k) - \varphi_\lambda(y^* + x, k)}{D(x, k, V_\lambda, W_\lambda)} \left[aV_\lambda + bW_\lambda - u(\varphi_\lambda(y^* + x, k), k) - \frac{1-\lambda}{\lambda} U(\varphi_\lambda(y^* + x, k), k) \right] \quad (77)$$

where (here φ_λ stands for $\varphi_\lambda(y^* + x, k)$)

$$\begin{aligned}D(x, k, V_\lambda, W_\lambda) &= (y^* + x) \left[(a^2 + b^2)V_\lambda + 2abW_\lambda - \delta_1 u(\sigma_\lambda(k), k) - \frac{(1-\lambda)\delta_2}{\lambda} U(\sigma_\lambda(k), k) \right] \\ &\quad - A(aV_\lambda + bW_\lambda - u(\varphi_\lambda, k) - \frac{1-\lambda}{\lambda} U(\varphi_\lambda, k)) \\ A(x, k, V_\lambda, W_\lambda) &= (f - \varphi_\lambda) \left[\left(u''_{11}(f(k), k) + \frac{1-\lambda}{\lambda} U''_{11}(f(k), k) \right) f' + u''_{12}(f(k), k) \right. \\ &\quad \left. + \frac{1-\lambda}{\lambda} U''_{12}(f(k), k) \right] + (y^* + x) f'(k) + u'_2(\varphi_\lambda, k) + \frac{1-\lambda}{\lambda} U'_2(\varphi_\lambda, k)\end{aligned}$$

Further more, we have

$$\frac{dV_\lambda}{dx} = \frac{dV_\lambda}{dk} \frac{dk}{dx} = (y^* + x) \frac{f(k) - \varphi_\lambda}{D(x, k, V_\lambda, W_\lambda)} \left[aV_\lambda + bW_\lambda - u(\varphi_\lambda, k) - \frac{1-\lambda}{\lambda} U(\varphi_\lambda, k) \right]. \quad (78)$$

Step 4: Rescaling the time

We introduce a new variable s such that $D(x, k, V_\lambda, W_\lambda) ds = dx$, then the system becomes

$$\begin{aligned}\frac{dx}{ds} &= D(x, k, V_\lambda, W_\lambda) \\ \frac{dk}{ds} &= (f(k) - \varphi_\lambda) \left[aV_\lambda + bW_\lambda - u(\varphi_\lambda(y^* + x, k), k) - \frac{1-\lambda}{\lambda} U(\varphi_\lambda(y^* + x, k), k) \right] \\ \frac{dV_\lambda}{ds} &= (y^* + x) (f(k) - \varphi_\lambda(y^* + x, k)) \left[aV_\lambda + bW_\lambda - u(\varphi_\lambda, k) - \frac{1-\lambda}{\lambda} U(\varphi_\lambda, k) \right]\end{aligned}$$

We now eliminate W_λ , to get an equation in (x, k, V) only. From the equation of $\mu(k) = F(x, k)$, we have

$$bW_\lambda(k) = F(y^*(k) + x, k) + u(f(k), k) + \frac{1-\lambda}{\lambda} U(f(k), k) - aV_\lambda(k)$$

The dynamics of $(x(s), k(s), V_\lambda(s))$ are given by:

$$\begin{aligned} \frac{dx}{ds} &= \tilde{D}(x, k, V_\lambda) \\ \frac{dk}{ds} &= (f(k) - \varphi_\lambda) [F(y^*(k) + x, k) + u(f(k), k) + \frac{1-\lambda}{\lambda}U(f(k), k) \\ &\quad - u(\varphi_\lambda(y^* + x, k), k) - \frac{1-\lambda}{\lambda}U(\varphi_\lambda(y^* + x, k), k)] \\ \frac{dV_\lambda}{ds} &= (y^* + x)(f(k) - \varphi_\lambda(y^* + x, k)) [F(y^*(k) + x, k) + u(f(k), k) + \frac{1-\lambda}{\lambda}U(f(k), k) \\ &\quad - u(\varphi_\lambda(y^* + x, k), k) - \frac{1-\lambda}{\lambda}U(\varphi_\lambda(y^* + x, k), k)] \end{aligned} \quad (79)$$

where $\tilde{D}(x, k, V_\lambda) = D(x, k, V_\lambda, W_\lambda)$. For \tilde{D} to be C^2 , we need f to be C^3 and u, U to be C^4 .

Step 5: Linearizing the system

As we already noted, if $k = k_\infty$, then $\mu(k_\infty) = 0$ and $F(y^*(k_\infty) + x, k_\infty) = \mu(k_\infty) = 0$ has only one solution $x = 0$. Set $v_\infty = V_\lambda(k_\infty)$. We consider the system near the point $(x, k, V_\lambda) = (0, k_\infty, v_\infty)$. For simplicity, we write

$$y_\infty = y^*(k_\infty) = u'_1(f(k_\infty), k_\infty) + \frac{1-\lambda}{\lambda}U'_1(f(k_\infty), k_\infty)$$

Computing the linearized system at $(0, k_\infty, v_\infty)$, we find:

$$\frac{d}{ds} \begin{pmatrix} x \\ k - k_\infty \\ V_\lambda - v_\infty \end{pmatrix} = \begin{pmatrix} a_\infty & b_\infty & c_\infty \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ k - k_\infty \\ V_\lambda - v_\infty \end{pmatrix}.$$

where:

$$\begin{aligned} a_\infty &= y^*(k_\infty)^2 \frac{\partial \varphi_\lambda}{\partial y}(y^*(k_\infty), k_\infty) (f'(k_\infty) - \bar{g}_\lambda(k_\infty)) \\ \bar{g}_\lambda(k_\infty) &= \frac{\lambda \delta_1 u'_1(f(k_\infty), k_\infty) + (1-\lambda) \delta_2 U'_1(f(k_\infty), k_\infty)}{\lambda u'_1(f(k_\infty), k_\infty) + (1-\lambda) U'_1(f(k_\infty), k_\infty)} \\ &\quad - \frac{\lambda u'_2(f(k_\infty), k_\infty) + (1-\lambda) U'_2(f(k_\infty), k_\infty)}{\lambda u'_1(f(k_\infty), k_\infty) + (1-\lambda) U'_1(f(k_\infty), k_\infty)} \end{aligned}$$

By the concavity assumptions, we find:

$$\begin{aligned} y_\infty &= u'_{1\infty} + \frac{1-\lambda}{\lambda}U'_{1\infty} = u'_1(f(k_\infty), k_\infty) + \frac{1-\lambda}{\lambda}U'_1(f(k_\infty), k_\infty) > 0, \\ u'_{11}(f(k_\infty), k_\infty) + \frac{1-\lambda}{\lambda}U'_{11}(f(k_\infty), k_\infty) &< 0 \\ \frac{\partial \varphi_\lambda}{\partial y}(y^*(k_\infty), k_\infty) &= \left(u''_1(f(k_\infty), k_\infty) + \frac{\lambda}{1-\lambda}U''_1(f(k_\infty), k_\infty) \right)^{-1} < 0 \end{aligned}$$

For any k_∞ satisfying (73) with $\bar{g}_\lambda(k_\infty) - f'(k_\infty) \neq 0$, we then have $a_\infty \neq 0$. Introducing:

$$\tilde{x} = x + \frac{b_\infty}{a_\infty}(k - k_\infty) + \frac{c_\infty}{a_\infty}(V_\lambda - v_\infty), \quad (80)$$

we transform the system (79) for $(\tilde{x}(s), k(s), V_\lambda(s))$. The linearization at the origin is given by:

$$\frac{d}{ds} \begin{pmatrix} \tilde{x} \\ k - k_\infty \\ V_\lambda - v_\infty \end{pmatrix} = \begin{pmatrix} -y_\infty^2 \varphi'_{1\infty} (\bar{g}_\lambda(k_\infty) - f'(k_\infty)) & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} \tilde{x} \\ k - k_\infty \\ V_\lambda - v_\infty \end{pmatrix}.$$

Step 6: Applying the center manifold theorem

By the center manifold theorem (cf. Theorem 1 of [8]), there exist an $\epsilon > 0$, and a map $h(k, V_\lambda)$, defined in a neighborhood $\mathcal{O} = (k_\infty - \epsilon, k_\infty + \epsilon) \times (v_\infty - \epsilon, v_\infty + \epsilon)$ of (k_∞, v_∞) such that

$$h(k_\infty, v_\infty) = 0, \quad \frac{\partial h}{\partial k}(k_\infty, v_\infty) = 0, \quad \frac{\partial h}{\partial V_\lambda}(k_\infty, v_\infty, 0) = 0$$

and the manifold \mathcal{M} defined by

$$\mathcal{M} = \left\{ \left(h(k, V_\lambda) - \frac{b_\infty}{a_\infty}(k - k_\infty) - \frac{c_\infty}{a_\infty}(V_\lambda - v_\infty), k, V_\lambda \right) \mid (k, V_\lambda) \in \mathcal{O} \right\},$$

is invariant under the flow associated to the system (79). The map h and the central manifold \mathcal{M} are C^2 , and \mathcal{M} is two-dimensional and tangent to the critical plane defined by $\tilde{x} = 0$.

If $k = k_\infty$ and $V_\lambda = v_\infty$, then $x = 0$, and $h(k_\infty, v_\infty) = \tilde{x}(k_\infty, v_\infty) = x(k_\infty, v_\infty) = 0$

We are interested in the solutions which lie on the central manifold \mathcal{M} . Writing:

$$x = h(k, V_\lambda) - \frac{b_\infty}{a_\infty}(k - k_\infty) - \frac{c_\infty}{a_\infty}(V_\lambda - v_\infty)$$

in the equation $\frac{dV}{dk} = y^*(k) + x$, we get

$$\begin{cases} \frac{dV_\lambda}{dk} &= u'_1(f(k), k) + \frac{1-\lambda}{\lambda} U'_1(f(k), k) + h(k, V_\lambda) - \frac{b_\infty}{a_\infty}(k - k_\infty) - \frac{c_\infty}{a_\infty}(V_\lambda - v_\infty), \\ V_\lambda(k_\infty) &= v_\infty \end{cases} \quad (81)$$

which can be viewed as eliminating the variable s from the second and third equations of the system (79). Since $a_\infty \neq 0$ and the right hand side of the first equation of (81) is continuously differentiable in $\mathcal{O}_1 = (k_\infty - \epsilon, k_\infty + \epsilon)$, therefore, is locally Lipschitz continuous. By $\frac{dV_\lambda}{dk}|_{k=k_\infty} = y_\infty \neq 0$ which follows from (76) and the first equation of (81), the nonconstant solution of this initial-value problem exist in \mathcal{O}_1 which we denote by $V_\lambda(k) = \zeta(k)$, where $\zeta(k_\infty) = v_\infty$ and $\zeta \in C^2(\mathcal{O}_1)$ if $h \in C^2(\mathcal{O})$. Substituting $V_\lambda(k) = \zeta(k)$ into $x = h(k, V_\lambda) - \frac{b_\infty}{a_\infty}(k - k_\infty) - \frac{c_\infty}{a_\infty}(V_\lambda - v_\infty)$ yields

$$x(k) = h(k, \zeta(k)) - \frac{b_\infty}{a_\infty}(k - k_\infty) - \frac{c_\infty}{a_\infty}(\zeta(k) - v_\infty).$$

Finally,

$$\mu(k) = F(u'_1(f(k), k) + \frac{1-\lambda}{\lambda} U'_1(f(k), k) + x(k), k)$$

and

$$W_\lambda(k) = \frac{1}{b}(\mu(k) + u(f(k), k) + \frac{1-\lambda}{\lambda} U(f(k), k) - a\zeta(k)$$

is also C^2 , so we have found a C^2 solution of the system (58),(59) with boundary conditions (62),(63).

4.3 The existence of equilibrium strategies

Introduce another function $\underline{g}_\lambda(k)$ defined by:

$$\underline{g}_\lambda(k) = \frac{\lambda u'_1(f(k), k) + (1-\lambda)U'_1(f(k), k)}{\frac{\lambda}{\delta_1} u'_1(f(k), k) + \frac{1-\lambda}{\delta_2} U'_1(f(k), k)} - \frac{\lambda \delta_2 u'_2(f(k), k) + (1-\lambda) \delta_1 U'_2(f(k), k)}{\lambda \delta_2 u'_1(f(k), k) + (1-\lambda) \delta_1 U'_1(f(k), k)} \quad (82)$$

Theorem 11 Suppose $f'(k_\infty)$ lies between $\underline{g}_\lambda(k_\infty)$ and $\bar{g}_\lambda(k_\infty)$. Then there exists an equilibrium strategy converging to k_∞

Comparing (72) and (82), we find that:

$$\bar{g}_\lambda(k) - \underline{g}_\lambda(k) = \lambda(1-\lambda)(\delta_1 - \delta_2) \frac{(\delta_1 - \delta_2) u'_1 U'_1 - u'_2 U'_1 + u'_1 U'_2}{(\lambda u'_1 + (1-\lambda) U'_1)(\lambda \delta_2 u'_1 + (1-\lambda) \delta_1 U'_1)}$$

So the sign of $\bar{g}_\lambda(k) - \underline{g}_\lambda(k)$ is the sign of $(\delta_1 - \delta_2) u'_1 U'_1 - u'_2 U'_1 + u'_1 U'_2$:

$$\begin{aligned} \delta_1 - \delta_2 > \frac{u'_2}{u'_1} - \frac{U'_2}{U'_1} &\implies \bar{g}_\lambda(k) > \underline{g}_\lambda(k) \\ \delta_1 - \delta_2 < \frac{u'_2}{u'_1} - \frac{U'_2}{U'_1} &\implies \bar{g}_\lambda(k) < \underline{g}_\lambda(k) \end{aligned}$$

Let us give some examples:

Example 1: $u(c) = U(c)$

In this case, $u = U$ and they do not depend on k . We get:

$$\begin{aligned} \bar{g}_\lambda(k) &= \lambda \delta_1 + (1-\lambda) \delta_2 \\ \underline{g}_\lambda(k) &= \frac{1}{\frac{\lambda}{\delta_1} + \frac{(1-\lambda)}{\delta_2}} \end{aligned}$$

These formulas do not depend on the utility function $u(c)$. They were first derived in [14] for the special case $u(c) = \ln c$. Note that $\lambda \delta_1 + (1-\lambda) \delta_2$ is the arithmetic mean, and $\left(\frac{\lambda}{\delta_1} + \frac{(1-\lambda)}{\delta_2}\right)^{-1}$ is the geometric mean.

Example 2: $u(c, k) = U(c, k)$

In that case, the criterion (55) becomes:

$$\int_0^\infty (\lambda e^{-\delta_1 t} + (1-\lambda) e^{-\delta_2 t}) u(c(t), k(t)) dt \quad (83)$$

This problem was studied in [14] for $u(c, k) = U(c, k) = \ln c$. In the case at hand, (83), we find:

$$\begin{aligned} \bar{g}_\lambda(k) &= \lambda \delta_1 + (1-\lambda) \delta_2 - \frac{u'_2(f(k), k)}{u'_1(f(k), k)} \\ \underline{g}_\lambda(k) &= \frac{1}{\frac{\lambda}{\delta_1} + \frac{(1-\lambda)}{\delta_2}} - \frac{u'_2(f(k), k)}{u'_1(f(k), k)} \end{aligned}$$

These are the same as the preceding ones, with the corrective term $-u'_2/u'_1$.

Example 3: $u = u(c)$ and $U = U(c)$

In that case, the criterion (55) becomes:

$$\lambda \int_0^\infty e^{-\delta_1 t} u(c(t)) dt + (1-\lambda) \int_0^\infty e^{-\delta_2 t} U(c(t)) dt$$

We find:

$$\begin{aligned} \bar{g}_\lambda(k) &= \frac{\lambda \delta_1 u'_1(f(k)) + (1-\lambda) \delta_2 U'_1(f(k))}{\lambda u'_1(f(k)) + (1-\lambda) U'_1(f(k))} \\ \underline{g}_\lambda(k) &= \frac{\lambda u'_1(f(k)) + (1-\lambda) U'_1(f(k))}{\frac{\lambda}{\delta_1} u'_1(f(k)) + \frac{(1-\lambda)}{\delta_2} U'_1(f(k))} \end{aligned}$$

Here again we find $\underline{g}_\lambda(k) < \bar{g}_\lambda(k)$.

We now proceed to the proof of Theorem 11. We apply Theorem 10, and we denote by V_λ and W_λ the solution of equations (58) and (59) with the boundary conditions (62) and (63). Set $\sigma_\lambda(k) := \varphi_\lambda(V'_\lambda(k), k)$, where φ_λ is defined by (60) or (61). We will now show that the corresponding trajectory $k(t)$, defined by (35), converges to k_∞ . By Proposition 8, this will prove that σ_λ is an equilibrium strategy.

We need the value of $V'_\lambda(k_\infty)$ and $\sigma'_\lambda(k_\infty)$. Differentiating (58), we find:

$$\begin{aligned} V'_\lambda(k_\infty)f(k_\infty) - V'_\lambda(k_\infty)\varphi_\lambda(V'_\lambda(k_\infty), k_\infty) + u(\varphi_\lambda(V'_\lambda(k_\infty), k_\infty), k_\infty) \\ + \frac{1-\lambda}{\lambda}U(\varphi_\lambda(V'_\lambda(k_\infty), k_\infty), k_\infty)) = u(f(k_\infty), k_\infty) + \frac{1-\lambda}{\lambda}U(f(k_\infty), k_\infty), \end{aligned}$$

thus $F(V'_\lambda(k_\infty), k_\infty) = 0$ and hence:

$$V'_\lambda(k_\infty) = y^*(k_\infty) = u'_1(f(k_\infty), k_\infty) + \frac{1-\lambda}{\lambda}U'_1(f(k_\infty), k_\infty). \quad (84)$$

To compute $\sigma'_\lambda(k_\infty)$, we consider (45), the integrated form of $V'_\lambda(k)$ where we substitute $h(t) = e^{-\delta t}$ and $H(t) = e^{-\delta_2 t}$, and replace U with $(1-\lambda)\lambda^{-1}$. Evaluating (45) at $k = k_\infty$, we get:

$$\begin{aligned} V'_\lambda(k_\infty) = \frac{u'_1(f(k_\infty), k_\infty)\sigma'_\lambda(k_\infty) + u'_2(f(k_\infty), k_\infty)}{\delta_1 - f'(k_\infty) + \sigma'_\lambda(k_\infty)} \\ + \frac{1-\lambda}{\lambda} \frac{U'_1(f(k_\infty), k_\infty)\sigma'_\lambda(k_\infty) + U'_2(f(k_\infty), k_\infty)}{\delta_2 - f'(k_\infty) + \sigma'_\lambda(k_\infty)}. \end{aligned} \quad (85)$$

Comparing (84) with (85), we can solve for $\sigma'_\lambda(k_\infty)$. We get:

$$\sigma'_\lambda(k_\infty) = f'(k_\infty) - \delta_1\delta_2 \frac{(\frac{1}{\delta_1}u'_1(f(k_\infty), k_\infty) + \frac{1-\lambda}{\lambda\delta_2}\delta U'_1(f(k_\infty), k_\infty))(g_\lambda(k_\infty) - f'(k_\infty))}{(u'_1(f(k_\infty), k_\infty) + \frac{1-\lambda}{\lambda}U'_1(f(k_\infty), k_\infty))(\bar{g}_\lambda(k_\infty) - f'(k_\infty))}, \quad (86)$$

Linearizing the equation of motion $\frac{dk}{dt} = f(k) - \sigma_\lambda(k)$ at $k = k_\infty$ yields

$$\frac{dk}{dt} = (f'(k_\infty) - \sigma'_\lambda(k_\infty))(k - k_\infty).$$

Then k converges to k_∞ if $f'(k_\infty) - \sigma'_\lambda(k_\infty) < 0$. Comparing with (86), and writing in the expressions of $\underline{g}_\lambda(k_\infty)$ and $\bar{g}_\lambda(k_\infty)$ yields:

$$\delta_1\delta_2 \frac{(\frac{1}{\delta_1}u'_1(f(k_\infty), k_\infty) + \frac{1-\lambda}{\lambda\delta_2}\delta U'_1(f(k_\infty), k_\infty))(g_\lambda(k_\infty) - f'(k_\infty))}{(u'_1(f(k_\infty), k_\infty) + \frac{1-\lambda}{\lambda}U'_1(f(k_\infty), k_\infty))(\bar{g}_\lambda(k_\infty) - f'(k_\infty))} < 0.$$

Because u and U are increasing with respect to their first variable, the first factor in the numerator and denominator are positive. We are left with:

$$\frac{\underline{g}_\lambda(k_\infty) - f'(k_\infty)}{\bar{g}_\lambda(k_\infty) - f'(k_\infty)} < 0$$

and the proof is complete.

5 The Chichilnisky criterion

In two influential papers [11], [10], Chichilnisky has proposed an axiomatic approach to sustainable development, based on the twin ideas that there should be no dictatorship of the present and no dictatorship of the future. She suggests to use the following criterion

$$I_\alpha(c(\cdot), k(\cdot)) := \delta \int_0^\infty u(c(t), k(t)) e^{-\delta t} dt + \alpha \lim_{t \rightarrow \infty} U(c(t), k(t)), \quad (87)$$

The coefficient $\delta > 0$ in front of the integral will make later formulas simpler.

Lemma 12 *Suppose $u(c, k) \geq 0$ and*

$$\sup \{U(c, k) \mid c > 0, k > 0, c = f(k)\} = \infty.$$

Then, for any $\alpha > 0$,

$$\sup \{I_\alpha(c(\cdot), k(\cdot)) \mid (c(\cdot), k(\cdot)) \in \mathcal{A}(k_0)\} = \infty,$$

where $\mathcal{A}(k_0)$ is defined below Definition 1.

Proof. For fixed $k_0 > 0$, pick any $A > 0$, and choose some constants (c_1, k_1) such that $c_1 = f(k_1)$ and $U(c_1, k_1) > A\alpha^{-1}$. Then choose some $c_0 \in (0, c_1)$ such that $f(k_0) - c_0 > 0$. With every $T > 0$, we associate the path $(c_T(t), k_T(t))$ defined by:

$$c_T(t) = \begin{cases} c_0 & \text{for } 0 \leq t \leq T, \\ c_1 & \text{for } T \leq t, \end{cases}$$

and denote the consumption-capital pair by (c_T, k_T) .

Since f is increasing, we have $\frac{dk}{dt} \geq f(k_0) - c_0$ for all t , so eventually $k(t)$ will reach the value k_1 . Choose for T the first time when $k(t) = k_1$. Writing this into the criterion, and remembering $c_1 = f(k_1)$, we find:

$$I_\alpha(c_T(\cdot), k_T(\cdot)) \geq \alpha \lim_{t \rightarrow \infty} U(c_T(t), k_T(t)) = \alpha U(c_1, k_1) \geq A.$$

■

This result shows that very often it is not possible to optimize I_α . Even when it is, there is the time-inconsistency problem: successive decision-makers will not agree on what the optimal solution is. This is seen most easily by considering the following criterion:

$$(1 - \alpha) \delta \int_0^\infty u(c(t), k(t)) e^{-\delta t} dt + \alpha r \int_0^\infty U(c(t), k(t)) e^{-rt} dt. \quad (88)$$

When $r \rightarrow 0$, the last term converges to $\alpha U(c_\infty, k_\infty)$, so that (88) converges to (87). On the other hand, Criterion (88) is a special case of the biexponential criterion (55) with $\delta_1 = \delta$, $\delta_2 = r$, and:

$$\frac{\alpha r}{(1 - \alpha) \delta} = \frac{1 - \lambda}{\lambda}, \quad \lambda = \frac{(1 - \alpha) \delta}{\alpha r + (1 - \alpha) \delta}, \quad 1 - \lambda = \frac{\alpha r}{\alpha r + (1 - \alpha) \delta} \quad (89)$$

So, for each $r > 0$, the criterion (88) gives rise to a time-inconsistent problem. Using the results in the preceding section, we find a continuum of equilibrium strategies. Substituting (89) into (72) and (82), we find the corresponding values:

$$\begin{aligned}\bar{g}_r(k) &= \frac{(1-\alpha)\delta^2 u'_1(f(k),k) + \alpha r^2 U'_1(f(k),k)}{(1-\alpha)\delta u'_1(f(k),k) + \alpha r U'_1(f(k),k)} - \frac{(1-\alpha)\delta u'_2(f(k),k) + \alpha r U'_2(f(k),k)}{(1-\alpha)\delta u'_1(f(k),k) + \alpha r U'_1(f(k),k)} \\ \underline{g}_r(k) &= \frac{(1-\alpha)\delta u'_1(f(k),k) + \alpha r U'_1(f(k),k)}{(1-\alpha)u'_1(f(k),k) + \alpha U'_1(f(k),k)} - \frac{(1-\alpha)u'_2(f(k),k) + \alpha U'_2(f(k),k)}{(1-\alpha)u'_1(f(k),k) + \alpha U'_1(f(k),k)}\end{aligned}$$

The equations (58), (59) become:

$$(f - \varphi_r(V'_r))V'_r + u(\varphi_r) + \frac{\alpha r}{(1-\alpha)\delta}U(V'_r) = \frac{\delta+r}{2}V_r + \frac{\delta-r}{2}W_r, \quad (90)$$

$$(f - \varphi_r(V'_r))W'_r + u(\varphi_r) + \frac{\alpha r}{(1-\alpha)\delta}U(V'_r) = \frac{\delta-r}{2}V_r + \frac{\delta+r}{2}W_r \quad (91)$$

where $\varphi_r(x, k)$ is defined by:

$$u'_1(\varphi_r(x, k), k) + \frac{\alpha r}{(1-\alpha)\delta}U'_1(\varphi_r(x, k), k) = x$$

and the boundary conditions (62), (63):

$$V_r(k_\infty) = \frac{1}{\delta}u(f(k_\infty), k_\infty) + \frac{\alpha}{(1-\alpha)\delta}U(f(k_\infty), k_\infty), \quad (92)$$

$$W_r(k_\infty) = \frac{1}{\delta}u(f(k_\infty), k_\infty) - \frac{\alpha}{(1-\alpha)\delta}U(f(k_\infty), k_\infty). \quad (93)$$

The equilibrium strategy is given by $\sigma_r(k) = \varphi_r(V'_r(k), k)$, and the corresponding trajectory converges to k_∞ . Note that these strategies are defined locally. More precisely, denote by $]k_r^-, k_r^+[$ the maximal interval of existence of the solution (V_r, W_r) of the ODE (90), (91) with the boundary condition (92), (93), so that $k_r^- < k_\infty < k_r^+$.

We shall now solve the Chichilnisky problem by setting $r = 0$ in the preceding equations. We have:

$$\bar{g}_0(k) = \delta - \frac{u'_2(f(k), k)}{u'_1(f(k), k)}, \quad (94)$$

$$\underline{g}_0(k) = \frac{(1-\alpha)\delta u'_1(f(k), k)}{(1-\alpha)u'_1(f(k), k) + \alpha U'_1(f(k), k)} - \frac{(1-\alpha)u'_2(f(k), k) + \alpha U'_2(f(k), k)}{(1-\alpha)u'_1(f(k), k) + \alpha U'_1(f(k), k)} \quad (95)$$

The pair (V_0, W_0) has to solve the following boundary-value problem:

$$(f(k) - \varphi_0(V'_0, k))V'_0 + u(\varphi_0(V'_0, k), k) = \frac{\delta}{2}(V_0 + W_0), \quad (96)$$

$$(f(k) - \varphi_0(V'_0, k))W'_0 + u(\varphi_0(V'_0, k), k) = \frac{\delta}{2}(V_0 + W_0), \quad (97)$$

$$V_0(k_\infty) = \frac{1}{\delta}u(f(k_\infty), k_\infty) + \frac{\alpha}{(1-\alpha)\delta}U(f(k_\infty), k_\infty), \quad (98)$$

$$W_0(k_\infty) = \frac{1}{\delta}u(f(k_\infty), k_\infty) - \frac{\alpha}{(1-\alpha)\delta}U(f(k_\infty), k_\infty) \quad (99)$$

with:

$$u'_1(\varphi_0(x, k), k) = x, \quad \varphi_0(u'_1(c, k), k) = c. \quad (100)$$

Subtracting (97) from (96), we get:

$$(f(k) - \varphi_0(V'_0, k))(V'_0 - W'_0) = 0.$$

Similarly to Lemma 9, it follows that $V'_0 - W'_0 = 0$, and so $V_0 - W_0$ is a constant, namely:

$$W_0(k) = V_0(k) + W_0(k_\infty) - V_0(k_\infty) = V_0(k) - 2\frac{\alpha}{(1-\alpha)\delta}U(f(k_\infty), k_\infty).$$

Writing this in (96) and (97), we find that $V_0(k)$ is a solution of the boundary-value problem:

$$\begin{aligned} (f(k) - \varphi_0(V'_0, k))V'_0 + u(\varphi_0(V'_0, k), k) &= \delta V_0 - \frac{\alpha}{1-\alpha}U(f(k_\infty), k_\infty), \\ V_0(k_\infty) &= \frac{1}{\delta}u(f(k_\infty), k_\infty) + \frac{\alpha}{(1-\alpha)\delta}U(f(k_\infty), k_\infty). \end{aligned}$$

Setting $V := V_0 - \frac{\alpha}{(1-\alpha)\delta}U(f(k_\infty), k_\infty)$, we see that V is a solution of the boundary-value problem:

$$(f(k) - \varphi_0(V', k))V' + u(\varphi_0(V', k), k) = \delta V, \quad (101)$$

$$V(k_\infty) = \frac{1}{\delta}u(f(k_\infty), k_\infty). \quad (102)$$

This problem has been studied in [15] (Section 2.4, case 2) when $u(c, k)$ does not depend on k . In the general case, we have:

Proposition 13 *If*

$$f'(k_\infty) \neq \delta - \frac{u'_2(f(k_\infty), k_\infty)}{u'_1(f(k_\infty), k_\infty)} \quad (103)$$

the problem (101), (102) has two solutions V_1 and V_2 , defined on some non-empty half-interval $[k_\infty, k_\infty + a)$ or $(k_\infty - a, k_\infty]$. Both are C^1 on the half-interval, C^2 on the interior, and have the same derivative at k_∞ , given by:

$$\varphi_0(V'_i(k_\infty), k_\infty) = c_\infty = f(k_\infty), \quad i = 1, 2.$$

Proof. From now on, write φ instead of φ_0 . Rewrite (101) as a Pfaff system:

$$dV = pdk, \quad (104)$$

$$p(f(k) - \varphi(p, k)) + u(\varphi(p, k), k) = \delta V \quad (105)$$

Differentiating (105) leads to:

$$\delta dV = (f(k) - \varphi(p, k))dp + pf'(k)dk + u'_2(\varphi(p, k), k)dk,$$

where we used (100). Together with (104), this yields

$$(f(k) - \varphi(p, k))dp = [\delta p - pf'(k) - u'_2(\varphi(p, k), k)]dk. \quad (106)$$

We have to investigate this system near the point $k = k_\infty$ and $V = V_\infty$. Writing (105) at this point, we get:

$$p(f(k_\infty) - \varphi(p, k_\infty)) + u(\varphi(p, k_\infty), k_\infty) = \delta V_\infty \quad (107)$$

which has to be solved for p . Note that, because of (100), we have:

$$\min_p \{(f(k_\infty) - \varphi(p, k_\infty))p + u(\varphi(p, k_\infty), k_\infty)\} = u(f(k_\infty), k_\infty) \quad (108)$$

In the case at hand, we have $\delta V_\infty = u(f(k_\infty), k_\infty)$, but for the sake of completeness, and to have a full description of the phase space in the (k, V) plane, we will first investigate the cases $\delta V_\infty < u(f(k_\infty), k_\infty)$ and $\delta V_\infty > u(f(k_\infty), k_\infty)$.

Case 0: $\delta V_\infty < u(f(k_\infty), k_\infty)$

Because of (108), equation (107) has no solution. So there are no solutions going through (k_∞, V_∞) .

Case 1. $\delta V_\infty > u(f(k_\infty), k_\infty)$.

Equation (107) has two distinct solutions $p_1 \neq p_2$. Note that neither p_1 nor p_2 minimize the left hand side so $f(k_\infty) - \varphi(p_i, k_\infty) \neq 0$ for $i = 1, 2$. We may therefore consider the initial value problem:

$$\frac{dp}{dk} = \frac{\delta p - pf'(k) - u'_2(\varphi(p, k), k)}{f(k) - \varphi(p, k)}, \quad p(k_\infty) = p_i.$$

It has a well-defined smooth solution $p_i(k)$, defined in a neighborhood of k_∞ . We then define a function V_i by:

$$V_i(k) := \frac{1}{\delta} [(f(k) - \varphi(p, k))p + u(\varphi(p, k), k)].$$

Notice that $V_i(k_\infty) = V_\infty$, so $V_i(k)$ solves the initial value problem (101), (102), with $V'_i(k_\infty) = p_i$. Taking $i = 1, 2$, we find two solutions $V_1(k)$ and $V_2(k)$ of the same initial value problem, with $V'_1(k_\infty) \neq V'_2(k_\infty)$.

Case 2. $\delta V_\infty = u(f(k_\infty), k_\infty)$ and $f'(k_\infty) \neq \delta - u'_2/u'_1$.

Equation (107) then has a single solution p_0 , and we have $f(k_\infty) = \varphi(p_0, k_\infty)$, so that:

$$p_0 = u'_1(\varphi(p_0, k_\infty), k_\infty) = u'_1(f(k_\infty), k_\infty) > 0.$$

In this case, we shall use the same system (104)-(105), but we will take p instead of k as the independent variable. We consider the initial value problem

$$\frac{dk}{dp} = \frac{f(k) - \varphi(p, k)}{p(\delta - f'(k) - \frac{u'_2(\varphi(p, k), k)}{p})}, \quad k(p_0) = k_\infty.$$

It has a C^2 solution $k(p)$, defined in a neighborhood of $p = p_0$. We associate with it a curve in the phase space (k, V) , defined in parametric form by the equations

$$\begin{aligned} k &= k(p), \\ V &= \frac{1}{\delta} [(f(k) - \varphi(p, k))p + u(\varphi(p, k), k)]. \end{aligned}$$

Then V is also C^2 with respect to p near $p = p_0$. Moreover, we have

$$\begin{aligned}\frac{dV}{dp} &= \frac{1}{\delta} [(f(k) - \varphi(p, k) - \varphi'_1(p, k)p + u'_1(\varphi(p, k), k)\varphi'_1(p, k))] \\ &\quad + \frac{1}{\delta} [(f'(k) - \varphi'_2(p, k))p + u'_1(\varphi(p, k), k)\varphi'_2(p, k) + u'_2(\varphi(p, k), k)] \frac{dk}{dp} \\ &= \frac{f(k) - \varphi(p, k)}{\delta - f'(k) - \frac{u'_2(\varphi(p, k), k)}{p}}\end{aligned}$$

Since $f(k_\infty) = \varphi(p_0, k_\infty)$, we obtain

$$\begin{aligned}\frac{dk}{dp}(p_0) &= 0, \\ \frac{dV}{dp}(p_0) &= 0.\end{aligned}$$

This shows that the parametric curve $p \rightarrow (k(p), V(p))$ in the (k, V) -plane has a cusp at (k_∞, V_∞) . To find the type of the cusp, we compute the second order derivatives with respect to p . We find:

$$\frac{d^2k}{dp^2}(p_0) = \frac{-\varphi'_1(p_0, k_\infty)}{p_0[\delta - f'(k_\infty) - \frac{u'_2(f(k_\infty), k_\infty)}{u'_1(f(k_\infty), k_\infty)}]} = \frac{-u''_{11}(f(k_\infty), k_\infty)}{u'_1(f(k_\infty), k_\infty)[\delta - f'(k_\infty) - \frac{u'_2(f(k_\infty), k_\infty)}{u'_1(f(k_\infty), k_\infty)}]} \neq 0, \quad (109)$$

$$\frac{d^2V}{dp^2}(p_0) = \frac{-\varphi'_1(p_0, k_\infty)}{\delta - f'(k_\infty) - \frac{u'_2(f(k_\infty), k_\infty)}{u'_1(f(k_\infty), k_\infty)}} = \frac{-u''_{11}(f(k_\infty), k_\infty)}{\delta - f'(k_\infty) - \frac{u'_2(f(k_\infty), k_\infty)}{u'_1(f(k_\infty), k_\infty)}} \neq 0, \quad (110)$$

This shows that the cusp has two branches, with a common tangent between them. The branches extend on the left of k_∞ if the left-hand side of (109) is negative, that is, if:

$$\delta - f'(k_\infty) - \frac{u'_2(f(k_\infty), k_\infty)}{u'_1(f(k_\infty), k_\infty)} < 0 \quad (111)$$

and on the right if it is positive. The slope m of the common tangent is given by

$$m = \frac{d^2V}{dp^2}(k_0) / \frac{d^2k}{dp^2}(p_0) = p_0,$$

and it is also the one-sided derivative of V at k_∞ . Note for future use that because $p_0 = u'_1(f(k_\infty), k_\infty)$ we have:

$$\frac{d}{dk} \frac{1}{\delta} u(f(k), k) \Big|_{k=k_\infty} = \frac{1}{\delta} (u'_1(f(k_\infty), k_\infty) f'(k_\infty) + u'_2(f(k_\infty), k_\infty)),$$

so that $m = p_0$ does not coincide with the tangent of the curve $\delta V = u(f(k))$ at $k = k_\infty$.

Case 3: $\delta V_\infty = u(f(k_\infty), k_\infty)$ and $f'(k_\infty) = \delta - u'_2/u'_1$

This is the case which was investigated in Theorem 3. We have shown that there exists a C^2 solution $V(k)$ on a neighbourhood of k_∞ .

This concludes the proof of Proposition 13 (in fact, we only need Case 2). ■

Let us summarize these results. The curve $\Gamma = \{(k, V) \mid \delta V = u(f(k), k)\}$ separates the plane in two regions.

- The region below the curve corresponds to Case 0: there are no solution there.
- The region above the curve corresponds to Case 1: through each point (k_∞, V_∞) there are two smooth solutions intersecting transversally.
- If (k_∞, V_∞) is on the curve, but $f'(k_\infty) + \frac{u'_2(f(k_\infty), k_\infty)}{u'_1(f(k_\infty), k_\infty)} \neq \delta$, we are in Case 2. There are two C^1 solutions defined only on one side of k_∞ . They are tangent at (k_∞, V_∞) , and transversal to Γ .
- If (k_∞, V_∞) is on the curve, and $f'(k_\infty) + \frac{u'_2(f(k_\infty), k_\infty)}{u'_1(f(k_\infty), k_\infty)} = \delta$, we are in Case 3: there is a C^2 solution defined on a neighbourhood of k_∞ .the

Figure 1 gives the phase diagram in the (k, V) plane:




Figure 1. The illustration of the solutions.

Proposition 13 gives us two solutions, $V_i, i = 1, 2$. Each of them gives rise to an strategy σ_i through the formula $\sigma_i(k) = \varphi_0(V'_i(k), k)$ with φ_0 defined by (100). The strategy σ_i is C^0 on the half-interval, C^1 on its interior, with $\sigma_i(k_\infty) = f(k_\infty)$.

Proposition 14 *One, and only one, of the strategies σ_1 and σ_2 , converges to k_∞ .*

Proof. Consider the Euler-Lagrange equation (16) for the Ramsey problem. The phase diagram is given in Figure 2, where \underline{k} is defined by $f'(\underline{k}) = \delta - u'_2/u'_1$.

If $k_\infty \neq \underline{k}$, there is one trajectory \mathcal{T} going through $(k_\infty, u(f(k_\infty), k_\infty))$. The point $(k_\infty, u(f(k_\infty), k_\infty))$ separates it into two branches, the upper one and the lower one. One of them goes to $(k_\infty, u(f(k_\infty), k_\infty))$, and the other one leaves it. These two branches are also the trajectories associated with the two strategies σ_1 and σ_2 , so one of them converges and the other diverges. ■

Neither depends on k , and we have:

$$\begin{aligned}\bar{g}_0(k) &= \delta, \\ \underline{g}_0(k) &= (1 - \alpha) \delta\end{aligned}$$

We have $\underline{g}_0(k) < \bar{g}_0(k)$, so the equilibrium strategy exists only on the right hand side of $\underline{g}_0(k)$. The existence condition is:

$$(1 - \alpha) \delta < f'(k_\infty) < \delta$$

and the equilibrium strategy σ is defined on $[k_\infty, \infty[$. We denote $f'^{-1}(\delta)$ and $f'^{-1}((1 - \alpha)\delta)$ by \underline{k} and \bar{k} respectively. There are three cases, depending on the position of the initial point k_0 :

- If $k_0 > \bar{k}$, then, for any $k_\infty \in]\bar{k}, k_0[$, there exists an equilibrium strategy starting from k_0 and converging to k_∞ .
- if $\underline{k} < k_0 < \bar{k}$, then, for any $k_\infty \in]\underline{k}, k_0[$, there exists an equilibrium strategy starting from k_0 and converging to k_∞ .
- if $k_0 < \underline{k}$, the only equilibrium strategy starting from k_0 is the optimal strategy for the Ramsey problem (that is, for the case $\alpha = 0$) which converges to the level \underline{k} where $f'(\underline{k}) = \delta$.

Example 2: $u(c, k) = U(c, k)$

We find:

$$\begin{aligned}\bar{g}_0(k) &= \delta - \frac{u'_2(f(k), k)}{u'_1(f(k), k)} \\ \underline{g}_0(k) &= (1 - \alpha) \delta - \frac{u'_2(f(k), k)}{u'_1(f(k), k)}\end{aligned}$$

The existence condition is:

$$(1 - \alpha) \delta - \frac{u'_2(f(k), k)}{u'_1(f(k), k)} < f'(k_\infty) < \delta - \frac{u'_2(f(k), k)}{u'_1(f(k), k)}$$

and the equilibrium strategy σ is defined on $[k_\infty, k_\infty + \kappa[$ for some $\kappa > 0$. The situation is similar to the preceding one, bearing in mind that now the strategy σ may be defined locally only

Example 3: $u = u(c)$ and $U = U(k)$

In that case, we find:

$$\begin{aligned}\bar{g}_0(k) &= \delta, \\ \underline{g}_0(k) &= \delta - \frac{\alpha}{1 - \alpha} \frac{U'_2(f(k), k)}{u'_1(f(k), k)}\end{aligned}$$

There are two subcases:

- if $\frac{U'_2(f(k), k)}{u'_1(f(k), k)} > 0$, then $\underline{g}_0(k) < \bar{g}_0(k)$. The equilibrium strategy then is defined on the right hand side of k_∞ , as in the preceding cases.

- [3] Geir Asheim, *Justifying, Characterizing and Indicating Sustainability*, Springer (2007) ISBN: 978-1-4020-6199-8
- [4] Martino Bardi and Italo Capuzzo-Dolcetta, *Optimal control and viscosity solutions of Hamilton-Jacobi-Bellman equations*, Springer Verlag, 1997
- [5] Robert Barro and Xavier Sala-i-Martin, *Economic growth*, MIT Press, 2003
- [6] Olivier J. Blanchard, Stanley Fischer, *Lectures on macroeconomics*, MIT Press. (1989).
- [7] Andrew Caplin, John Leahy, *The recursive approach to time inconsistency*, Journal of Economic Theory. (2006)131:134-156.
- [8] Jack Carr, *Applications of centre manifold theory*, Springer-Verlag. (1981).
- [9] David Cass, *Optimal growth in an aggregative model of capital accumulation*, Review of Economic Studies (1965)32: 233-240
- [10] Graciela Chichilnisky, *An axiomatic approach to sustainable development*, Social Choice and Welfare. (1996)13: 231-257.
- [11] Graciela Chichilnisky, *What is sustainable development?* Land Economics. (1997)4: 467-491.
- [12] Ivar Ekeland, Ali Lazrak, *Equilibrium policies when preferences are time-inconsistent*, <http://arxiv.org/abs/math/0808.3790>.(2006).
- [13] Ivar Ekeland, Ali Lazrak, *Being serious about non-commitment: subgame perfect equilibrium in continuous time*, <http://arxiv.org/abs/math/0604264>.(2006).
- [14] Ivar Ekeland, Ali Lazrak, *The golden rule when preferences are time-inconsistent*, Mathematics and Financial Economics. (2010)4:29-55.
- [15] Ivar Ekeland, *From Frank Ramsey to René Thom: a classical problem in the calculus of variations leading to an implicit differential equation*, Discrete and Continuous Dynamical Systems, 28, 3 (2010) p. 1101 - 1119
- [16] Ivar Ekeland, Larry Karp, Rashid Sumaila, *Equilibrium management of fisheries with overlapping altruistic generations*, Working paper (2011), <http://www.ceremade.dauphine.fr/~ekeland/Articles/Karp.pdf>
- [17] Ivar Ekeland, Roger Temam, *Convex analysis and variational problems*, SIAM Classics in Applied Mathematics
- [18] Frederick, Shane, Loewenstein, George, and O'Donoghue, Ted "Time discounting and time preference: a critical review", Journal of Economic Literature 40 (2002), 351-401
- [19] Hartman, *Ordinary differential equations*, Walter de Gruyter. (1990).
- [20] Geoffrey Heal, *Valuing the future: economic theory and sustainability*, Columbia University Press (1998)

- [21] Geoffrey Heal, *Intertemporal welfare economics and the environment*, in "Handbook of Environmental Economics", Volume 3, K.-G. Maler and J.R. Vincent ed., Elsevier (2005)
- [22] Larry Karp, *Non-constant discounting in continuous time*, Journal of Economic Theory. (2007)132:577-568
- [23] Larry Karp, In Ho Lee, *Time-consistent policies*, Journal of Economic Theory. (2003)112:353-364.
- [24] P. Krusell, A. Smith, *Consumption-savings decisions with quasi-geometric discounting*. Econometrica (2003)71:365–375
- [25] Tjalling Koopmans, *On the concept of optimal economic growth*, in "The economic approach to development planning", Elsevier (1965)
- [26] Yuri A.Kuznetsov, *Elements of applied bifurcation theory*, Springer (1998).
- [27] C. Harris, D.Laibson, *Dynamic choices of hyperbolic consumers*. Econometrica (2001)69:935–957Sdd
- [28] Chuan-Zhong Li, Karl-Gustav Löfgren, *Renewable resources and economic sustainability: a dynamic analysis with heterogeneous time preferences*. Journal of Environmental Economics and Management. (2000)40:236-250.
- [29] Edmund Phelps, R.A. Pollak, *On second-best national saving and game-equilibrium growth*. Rev. Econ.Stud.35 (1968), 185–199
- [30] Edmund Phelps, *The indeterminacy of game-equilibrium growth*. In: Phelps, E.S. (ed.) "Altruism, Morality and Economic theory", pp. 87–105. Russell Sage Foundation, New York (1975)
- [31] Frank P.Ramsey, *A mathematical theory of saving*, The Economic Journal. (1928)38:543-559.
- [32] David Romer, *Advanced macroeconomics*, McGraw-Hill
- [33] Ussif R.Sumaila, Carl J.Walters, *Intergenerational discounting: a new intuitive approach*, Ecological Economics (2005)52:135-142.